

Production of Tannase enzyme from Industrial Waste to abate Environmental Pollution

D.S.R.S.Prakash and M.Kiran Deedi

¹Department of Biotechnology, Adikavi Nannaya University, Rajamahendravaram, East Godavari District, Andhra Pradesh, INDIA.

²Department of Biochemistry, GSL Medical College, Rajamahendravaram, East Godavari District, Andhra Pradesh, INDIA.

Submission Date: 16-01-2020; Revision Date: 18-03-2020; Accepted Date: 12-04-2020

ABSTRACT

Enzymes and micro-organisms can be used for the bioconversion of biological materials into useful products. A few reports are available in the literature on the production of tannase through biotechnological routes. A novel route for the production of enzyme tannase using the principles of fermentation technology and organic chemistry. Biological degradation is adjudged as an important mechanism of organic chemical removal in natural systems, owing to its environmental compatibility and plasticity. Consequently, microbial metabolic potentials are widely exploited for a number of industrial applications including decommissioning of environmental pollutants. An important enzyme having various industrial applications is tannase (tannin acyl hydrolase). It is an extracellular enzyme falling under the hydrolase group of enzymes. Tannin acyl hydrolase (E.C.3.1.1.20) commonly called tannase, catalyzes the hydrolysis of ester and depside bonds in such hydrolysable tannins as tannic acid, thereby, releasing glucose and gallic acid. Tannase is an inducible enzyme produced only in the presence of tannic acid, or its end-product, i.e., gallic acid. The work deals with the extraction of tannins and production of tannase enzyme from Cashew testa waste by submerged fermentation using immobilized cells of mutated strain Cashew, *Anacardium occidentale*, is widely cultivated in India especially in coastal Andhra Pradesh. The Cashew testa after taking out the raw nuts has been thrown by cashew industries. The testa of Cashew has a long number of tannins. This disposed testa containing toxic chemicals like tannic acid from tannins causing environmental pollution by contaminating the local ground water.

Key words: Tannins, Tannase, Cashew husk, Enzyme extraction.

Correspondence:

DSRS Prakash,

Department of
Biotechnology,
Adikavi Nannaya
University,
Rajamahendravaram,
Andhra Pradesh, INDIA.
Phone no: +91 9291011999

Email: drdevaprakash@
gmail.com

INTRODUCTION

The selection of a substrate for large-scale enzyme production by fermentation depends upon its availability and cost. In this regard the waste residue of cashew husk was used as substrate for obtaining the desired fermented product. Tannins are water soluble polyphenolic compounds found in plants as secondary metabolites which can be grouped as hydrolysable and non-hydrolysable tannins. The presence of these substances

in the husk of cashew was initially extracted and their crude extract was used separately as a substrate for the production of tannase by *Aspergillus oryzae*. Tannase production as well as the biodegradation of the substrate reached maximum within 24 to 36 h against crude tannin extract obtained from *Anacardium occidentale*. Tannase production was higher by almost two fold in the presence of crude tannin compared to pure tannic acid used as a substrate. It seems that the industrial production of tannase, using husk extract of *Anacardium occidentale* can be a very simple and suitable alternative to presently used procedures. This method appears to be much more accurate than those reported earlier.

Microbial production of tannin acyl hydrolase (EC 3.1.1.20), commonly referred as tannase is well documented (Madhevan *et al.*)^[1] It is used widely in the manufacture of instant tea, acorn wine and gallic acid

SCAN QR CODE TO VIEW ONLINE



www.ajbls.org

DOI :
10.5530/ajbls.2020.9.10

(Lekha and Lonsane, 1997).^[2] Gallic acid is an important substrate for the synthesis of propyl gallate in the food industry and trimethoprim in the pharmaceutical industry (Hadi *et al.* 1994).^[3] Tannase also has potential applications in the clarification of bear and fruit juices, manufacture of coffee flavored soft drinks, improvement in the flavor of grape wine and as an analytical probe for determining the structures of naturally occurring gallic acid esters.

The industrial applications of tannase have not been fully exploited because of its high cost, although there are a large number of reports on the production of tannase by submerged fermentation. Most of these do not involve the identification of critical parameters for enzyme biosynthesis and their optimization. This enzyme is synthesized by a number of fungi. Based on a preliminary screening of various isolates and other available fungal cultures, one of these organisms, *Aspergillus oryzae*, was selected for further studies. The present studies were aimed at identifying key environmental parameters that play important roles in enzyme synthesis by *Aspergillus oryzae* and their optimization by submerged fermentation.

Most of the reported tannase-producing organisms are fungi (Aoki *et al.* 1976;^[4] Bhat *et al.* 1998;^[5] Mondal *et al.* 2001c;^[6] Ramirez-Coronel *et al.* 2003^[7] and only a few are bacteria (Deschamps *et al.* 1983;^[8] Mondal and Pati, 2000;^[9] Mondal *et al.* 2001a.^[10] Many authors studied tannase production by these organisms in the medium containing pure tannic acid acting as both inducer as well as available carbon source. Pure tannic acid is a very costly substrate and is not suitable for large-scale production of the enzyme. In this respect crude tannin could be cost effective and suitable for the commercial production of the enzyme. Agro-residues and forest products are generally considered the best source of tannin-rich substrate (Pandey *et al.* 1999).^[11] Production of tannase by *Rhizopus oryzae* and *Aspergillus foetidus* from the powdered fruits of Terminalia chebula and Caesalpinia digyna has been reported (Mukherjee and Banerjee, 2004).^[12] In this regard there are no reports on bacterial tannase production using tannin containing any agro-based substrates. In the present study we are reporting for the first time the production of tannase by *Aspergillus oryzae* through submerged fermentation of crude tannin extracted from the cashew husk.

MATERIALS AND METHODS

Micro-organism: The non-pathogenic tannase – producing strain of *Aspergillus oryzae* obtained from the NCIM Pune, was used in the present study.

PREPARATION OF INOCULUMS

Tannase being an adaptive enzyme, pre-Induced inoculum is required to be prepared. Inoculum was prepared by growing a loopful amount of stock culture of the Fungi in 50 ml sterile modified Czapek's dox medium and slightly modified mildew test (MT) medium is used. Medium will be autoclaved (21°C) for 20 min and allowed to cool at room temperature. The filter sterilized, tannic acid (2%) will be added to each sample. Composition of alternative medium was (g %, w/v): KH_2PO_4 , 5; NH_4NO_3 , 10; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1; $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$, 0.1; $\text{MnCl}_2 \cdot 6\text{H}_2\text{O}$, 0.02; $\text{NaMO}_4 \cdot 2\text{H}_2\text{O}$, 0.01; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.125 glucose, 25.

EXTRACTION OF CRUDE TANNINS

Collected cashew husk from cashew industries near to Visakhapatnam, were grained into small particles and dried in hot air oven at 60°C for 24 hr. The testa (50 g) were then mixed with distilled water (200 ml) and kept at room temperature overnight. After soaking, the mixture was autoclaved under pressure autoclaving for 30 min. The filtered solutions were used as source of crude tannin.

Detection of tannin by Thin layer chromatography

In this study, the detection of tannins by thin layer chromatography of cashew testa extract were identified. The results, showed that when the extract was run with two dimensional TLC, it gives three fractions, (Figure 1).




Figure 1: TLC ethanol extract from cashew husk (1-dry and milled: 2-dry and entire: 3-natural and milled: 4-natural and entire: A-tannic acid standard).

To obtain pure compounds, the appropriate chromatographic techniques were used. Followed by reverse-phase chromatography (in water and alcohol, or water, alcohol and acetone mixtures)

MEASUREMENT OF TANNIN BIODEGRADATION

The tannin content of the crude cashew testa extract was measured before and after fermentation by Folin – Denis method. The crude extract (0.2 ml) was initially diluted with 8.3 ml of distilled water and then mixed with 0.5 ml of Folin – Denis reagent. After proper mixing, 1 ml of 15% (w/v) Na_2CO_3 was added to it and kept in the dark for 30 min at room temperature. The absorbency of tannin was measured spectrophotometrically at 700 nm and its concentration was calculated using pure tannic acid as standard.

FERMENTATION PROCESS AND EXTRACTION OF TANNASE ENZYME

Tannase production by *Aspergillus oryzae* was achieved through submerged fermentation of crude tannin at 35°C in a rotary shaker (160 rpm). Different concentrations of tannin were prepared by diluting the measured crude tannin with distilled water. The pH of the medium was adjusted to 5.5 after sterilization. Fermentations were carried out separately in individual 250 ml Erlenmeyer flasks containing 50 ml modified Czapek's dox medium with (1% v/v) fresh inoculum. The cell – free fermented broth was used as the source of the enzyme. The growth of organism in culture media was monitored by measuring dry weight of the biomass (mg/ml).

ASSAY OF TANNASE

Tannase activity in the fermented medium was determined by the colorimetric method of Mondal *et al.* (2001b).^[13] For assay, 0.1 ml of enzyme was incubated with 0.3 ml of substrate tannic acid (1.0% w/v in 0.2 M citrate buffer, pH 5.0) at 50°C for 30 min. This reaction was terminated by the addition of 3 ml BSA solution (1 mg/ml), which also precipitated the residual tannic acid. A control reaction was done side by using heat-denatured enzyme. The tubes were then centrifuged (5000 x g, 10 min) and precipitate was dissolved in 2 ml of SDS-triethanolamine (1% w/v, SDS in 5% v/v, triethanolamine) solution. Absorbency was measured at 530 nm after addition of 1 ml of FeCl_3 (0.13 M).

The specific extinction co-efficient of tannic acid at 530 nm was 0.577. Using this co-efficient, one unit of

tannase activity is defined as the amount of enzyme required to hydrolyze 1 mm substrate (tannic acid) in 1 min at 50°C and pH 5.0.

RESULTS AND DISCUSSION

RESULTS

The selection of a substrate for large – scale enzyme production by fermentation depends upon its availability and cost. In this regard the waste residue of cashew husk was used as substrate for obtaining the desired fermented product (Pandey, *et al.* 1999).^[11]

Tannin contents of the bark of some commonly available plants were initially examined by paper chromatography and quantified by colorimetric method. Among the eight plant species tested, the maximum amount of tannin was found in the extract of the husk of cashew. Tannase production by *A. Oryzae* was studied using cashew testa tannins as submerged fermentation media. It has been found that extract of *A. occidentale* was the best for induction of tannase production. A similar type of timer related enzyme production by the same organism was also reported with pure tannic acid as substrate. Tannase production by the organism was found to be maximal in the extract of *A. occidentale*. It is not clear to us why the production of enzyme in the extract of *A. occidentale* is high, but we assume that there may be some inducing factors that accelerate enzyme synthesis.

In the present work, studies on the tannase production from *Aspergillus oryzae* using cashew husk was carried out and the results were given in the tables and figs. The effect of some parameters at different ranges was studied and their influence on the production was discussed in this paper.

OPTIMIZATION OF SOME PARAMETERS FOR MAXIMUM TANNASE PRODUCTION

EFFECT OF INCUBATION PERIOD ON TANNASE PRODUCTION

Table 1: Effect of Incubation period on Tannase production.

S.No.	Incubation period (h)	Tannase activity U/ml
1.	12	22.62
2.	24	30.62
3.	36	28.54
4.	48	25.42
5.	60	24.83
6.	72	20.62




Figure 1: Effect of Incubation period on Tannase production.

EFFECT OF PH ON TANNASE PRODUCTION

Table 2: Effect of pH on Tannase production.

S.No	pH	Enzyme activity (U/ml)
1	3.5	24.32
2	4	28.94
3	4.5	30.56
4	5	32.62
5	5.5	29.62
6	6	28.62




Figure 2: Effect of pH on Tannase Production.

DISCUSSION

The production of tannase has increased with increase in incubation time up to 48 hrs, with further increase in the incubation time, decrease in the tannase production was observed. This may be due to the starting of the declining phase of the organism after 48 hr. (Figure 1) shows that, an incubation period of 48 hrs was optimum for tannase production by *Aspergillus oryzae*. An optimum incubation period around 36 h has been reported for tannase activity in case of *A. aculeatus* D B F 9.

It could be concluded from the results that tannase from *Aspergillus oryzae* needed an acidic environment to be active. Fungal tannase is an acidic protein in general.

EFFECT OF TEMPERATURE ON TANNASE PRODUCTION

Table 3: Effect of Temperature on Tannase production.

S.No.	Temperature °C	Tannase activity U/ml
1.	25	22.20
2.	30	24.52
3.	35	26.85
4.	40	30.6
5.	45	28.12
6.	50	23.23




Figure 3: Effect of Temperature on Tannase production.

To Study the effect of initial pH on tannase production, the pH of the medium was varied from 3.5 – 6.0 using 1 N HCl and 1 N NaOH and fermentation was done as usual. The enzyme was active at acidic pH and activity decreased as the pH 5.0 (Figure 2). Maximum tannase activity was 32.62 U/ml by the *Aspergillus oryzae* after the optimum pH of 5.0. It could be concluded from the results that tannase from the *Aspergillus Oryzae* needed an acidic environment to be active. Fungal tannase is an acidic protein in general. The obtained results were tabulated in Table 2 and also shown in Figure 2. There are reports describing of the optimum pH as 5.5 in case of tannase obtained from *A. Oryzae*, (yamada H, *et al.* 1968)^[14] and 6.0 in case of tannase obtained from *P.chrysogenum* and *A. niger*, Barthony C, *et al.* (1994).^[15] To Study the effect of different temperatures on tannase production, the flasks containing medium kept at temperature range was varied from 25°C – 50°C. With a rise in temperature, the tannase production increased and optimum activity 30.62U/ml was recorded at 40°C (Figure 3). With a further increase in temperature, there was a decrease in activity. The optimum temperature for tannase production was 40°C. The obtained results

Table 4: Comparative study of tannase production in pure tannic acid and crude tannin extract as substrates by *Aspergillus oryzae*.

Substrates in fermentation medium	Growth (mg/ml)	Tannase (U/ml)
Crude tannin (0.5 g/100 ml)	0.82 ± 0.18	0.17 ± 0.06
Crude tannin (0.5 g/100 ml) + Basal salts*	1.12 ± 0.22	0.66 ± 0.12
Tannic acid (1.5 g/100 ml) containing enriched medium**	0.58 ± 0.12	0.31 ± 0.13

were tabulated in Table 3 and also shown in Figure 3. An optimum temperature around 30°C has been reported for tannase activity in case of *A. oryzae* (Iibuchi's, *et al.* 1968)^[16] and *P. chrysogenum*, around 35°C in case of *A. Niger* and 50°C in case of candida sp. (Aokoki *et al.* 1976).

Enzyme production was also studied at different concentration (0.5 – 2.0%, w/v) of crude tannins. It was observed that a specific concentration of crude tannin from a plant influenced enzyme production the strongest (Table 4).

In our experiments a maximal amount of enzyme was produced in medium containing 0.5% (w/v) of crude tannin of *A. occidentale*, but the highest enzyme production was observed when 1.5% (w/v) of pure tannic acid was used as substrate in the medium the concentration of tannin is thus a very important determining factor for tannase biosynthesis for most fungi and bacteria. The actual mode of tannase induction in a particular concentration of tannin has not been properly explained until now. Lewis and Starkey (1969)^[17] mentioned that higher concentrations of tannin lead to non-reversible bonds with surface proteins and impair the metabolism as well as growth of the organism. One of the most striking observations in this experiment is that enzyme production was increased about four-fold in the medium containing basal salt with crude tannin (0.5%) compared to medium containing crude tannin extract (0.5%) of *A. occidentale* alone (Table 1). This result revealed that some specific microelements (salts and ions) are probably essential for growth as well as enzyme synthesis by *A. Oryzae*. Both growth of the organism and enzyme production increased two-fold when it was grown in salt containing crude tannin extract rather than enriched pure tannic acid medium. All these beneficial effects of the plant extract of *A. occidentale* make it promising as one of the best as well as cheaper substrates for the large scale production of microbial tannase.

CONCLUSION

Tannase has now been extensively used in different biochemical industries. The selected bacterium used in this study is able to synthesize high amounts of tannase through fermentation of crude tannin of *A. Occidentale*. Exploitation of these plant extracts could be a source of cheaper substrate for industrial production of microbial tannase.

ACKNOWLEDGEMENT

Authors express their sincere gratitude for giving support to publish this paper to Prof. M. Jagannadha Rao, Vice Chancellor, Adikavi Nannaya University, Rajamahendravaram, East Godavari District, Andhra Pradesh, India.

CONFLICT OF INTEREST

Non declared

ABBREVIATIONS

NCIM: National Collection of Industrial Microorganisms; **MT:** Mildew Test; **TLC:** Thin Layer Chromatography; **SDS:** Sodium dodecyl sulfate.

REFERENCES

- Mahadevan A, Sivasweami S. In Frontiers in applied microbiology. Print-house, India. 1985;327.
- Lekha PK, Lonsane BK. Production and application of Tannin acyl hydrolase. *Advances in Applied Microbiology*. 1997;44:215-60.
- Hadi TAR, Banerjee BCB. Optimization of tannase biosynthesis by anewly isolated *Rhizopus oryzae*. *Bioprocess Eng*. 1994;11:239-43.
- Aoki K, Shinke R, Nishira H. Purification and some properties of yeast tannase. *Agricultural and Biological Chemistry*. 1976;40(1):79-85.
- Bhat TH, Singh B, Sharma OP. Microbial degradation of tannins: A current perspective. *Biodegradation*. 1998;9(5):343-57.
- Mondal HC, Banerjee D, Banerjee R, Pati BR. Production and Characterization of tannase from *Bacillus cereus* HBR6. *J Gen Appl Microbiol*. 2001a;47(5):263-7.
- Ramirez-Coronel A, Darvill A, Viniagra-Gonzalez G, Augur C. A novel tannase from *Aspergillus niger* with β -glucosidase activity. *Microbiology*. 2003;149(10): 2941-6.
- Deschamps AM, Otuk G, MLebeault J. Production of tannase and degradation of chestnut tannin by bacteria. *J Ferment Technol*. 1983;61(1):55-9.
- Mondal HC, Samanta S, Giri S, Pati BR. Distribution of tannic acid degrading microorganisms in the soil and comparative study of tannase from two fungal strains. *Acta Microbiol Pol*. 2001c;50(1):75-82.
- Mondal HC, Banerjee R, Pati BR. Tannase production by *Bacillus licheniformis*. *Biotechnol Lett*. 2000;22(9):767-9.
- Pandey A, Selvakumar P, Soccol CR, Nigam P. Solid-state fermentation for the production of industrial enzymes. *Curr Sci*. 1999;77:149-62.
- Mukherjee G, Banerjee R. Biosynthesis of tannase and gallic acid from tannin rich substrates by *Rhizopus oryzae* and *Aspergillus foetidus*. *J Basic Microbiol*. 2004;44(1):42-8.
- Mondal HC, Banerjee D, Jana M, Pati BR. Colorimetric assay method for determination of the tannin acyl hydrolase (EC 3.1.1.20) activity. *Anal Biochem*. 2001b;295(2):168-71.

14. Ibuchi S, Minoda Y, Yamada K. Studies on tannin acyl hydrolase of the microorganisms. Part III. Purification of the enzyme and some properties of it. *Agricultural and Biological Chemistry*. 1950;32(7):803-9.
15. Barthony C, Regeat F, Pourrat H. Production, purification and characterization of a tannase from *Aspergillus niger* LCF 8. *Journal of Fermentation Technology*. 1994;77(3):320-3.
16. libuchi S, Minoda Y, Yamada K. A new method of determining the enzyme activity using change in the ultraviolet absorption. *Agricultural and Biological Chemistry*. 1968;31(5):513-8.
17. Lewis JA, Starkey RL. Decomposition of plant tannins by some soil microorganisms. *Soil Science*. 1969;107(4):235-41.

Cite this article: Prakash DSRS, Deedi MK. Tannase enzyme Production from Industrial Waste to abate Environmental Pollution. *Asian J Biol Life Sci*. 2020;9(1):68-73.