

Comparative Phytochemical Analysis of Two Wound Healing Flora

Sreeranjini Kundamveettil

Post-Graduate Department of Botany, Little Flower College (Affiliation to University of Calicut), Guruvayur, Kerala, INDIA.

Submission Date: 05-03-2024; Revision Date: 19-03-2024; Accepted Date: 10-04-2024.

ABSTRACT

Chromolaena odorata (L.) R.M. King & H. Rob., shrub of Family Asteraceae and *Hemigraphis alternata* (Burm.f.) T. Anderson, herb of Family Acanthaceae are widely used as wound healers in local medicine. The leaf extract of both plants is used in local medicine which immediately prevent bleeding from wounds. The two plants possess several phytochemicals which can be cited as responsible for the wound healing process, but the presence of common phytochemicals in these two plants have not been worked out. Leaf extract studied using Gas Chromatography- Mass spectrometry method and phytochemicals identified by mass spectra. The comparative study of the components revealed the presence of common compounds including steroids, phthalates and sesquiterpenes - Diazoprogestosterone, Diisooctyl phthalate and Caryophyllene derivatives causing blood clotting, wound healing, preventing inflammations and pathogenic attack. The presence of terpenes can be concluded as reason for the wound healing property as they have antioxidant and anti-inflammatory action; presence of steroids may be accounted for the blood clotting process and the presence of phthalates considered as preventing infection in the wound areas. Isolation and optimization of components can help in modern medical practices instead of chemical combinations.

Keywords: *Chromolaena odorata*, *Hemigraphis alternata*, GC-MS, Sesquiterpenes, Phthalates.

Correspondence:

Dr. Sreeranjini Kundamveettil
Post Graduate
Department of Botany,
Little Flower College,
(Affiliation to University
of Calicut), Guruvayur,
Kerala, INDIA.

Email:
ranjini12000@gmail.com

INTRODUCTION

Unveiling of ethnomedicinal efficacies of flora has always been a boon to modern medicine. Human civilizations had utilised the healing powers of plants throughout the evolutionary period in different cultures. Majority drugs used in modern medicine claim their sources from plants. But along with this the inclusion of synthetic drugs is creating life-long complications in human life. Understanding this potential danger, retrospective outlook to nature has commenced. Drug development from plants can be factualised by intriguing to the traditional use of plant, identifying the

chemical content, conducting toxicity analysis of the components, randomised selection of one or several properties. Ethnopharmacology has been a corner stone for the entire procedure. In conjuncture with this basic idea, two plants *Chromolaena odorata* and *Hemigraphis alternata*, two wound healing ethnic plants were used for phytochemical studies for the constituents helpful in wound healing property. *Chromolaena odorata* (L.) R.M. King and H. Rob., a perennial shrub belonging to family Asteraceae, identified much for its wound healing as well as invasive properties is native to America.^[1] The plant is used for treating cuts, infections and various ailments. *Chromolaena odorata* is reported to possess several medically relevant properties.^[2] Reports are there on the detection of tannins, steroids, phenolics, flavonoids, saponins and terpenoids in the plant leaves.^[2] *Hemigraphis alternata* (Burm.f.) T. Anderson, a herbaceous perennial with a creeping growth habit is widely used in Kerala for its incredible potency to heal wounds. Apart from wound healing, the screening of various bioactive compounds from plant has led to the discovery of

SCAN QR CODE TO VIEW ONLINE



www.ajbls.com

DOI: 10.5530/ajbls.2023.13.29

anti-inflammatory, antidiuretic, anti-diabetic, antiulcer activity.^[3] The common wound healing property of the plant can be attributed to some bioactive compounds of the plants. Since the plant leaves are used in crude form for traditional wound healing, the leaves are subjected to phytochemical analysis. Biological relevance of the plant can be identified by basic pharmacological and chemical properties of plant by phytochemical screening methods like spectrometry and chromatography.^[4] Bioactive compounds and functional groups in plants are usually tracked by Fourier-Transform Infrared (FTIR) and gas chromatography-mass spectrometry (GC-MS).^[5] In this study, therefore a comparative analysis of the phytocomponents observed in both plants are done to elucidate the wound healing property.

MATERIALS AND METHODS


The leaves of *Chromolaena odorata* and *Hemigraphis alternata* was collected, cleaned, shade dried and powdered. The powder obtained was weighed out and 75 g of each of this was mixed with 150 mL of ethyl acetate and placed in the rotary shaker for 24 hr to separate the components. After 24 hr, the extract was filtered and the filtrate was allowed to dry out at room temperature for 2 days in dark. The extract (2 mL) was taken and phytochemical components were identified using GC-MS method at KFRI, Peechi. The specifications of the instrument used was Shimadzu GC-MS; Model Number: QP2010S; Column ELITE-5MS; 30 meter length; 0.25 mm ID; 0.25 μ m thickness. GCMS Software: GCMS Solutions and Libraries used:

NIST 11 and WILEY 8. A comparison of the GC-MS profile was done for the similar components.

RESULTS


2 mL extract of each plant was used for phytochemical analysis. The compounds identified in *Chromolaena odorata* are Cyclohexene, 5,6-diethenyl-1-methyl-, Cyclohexene, 3,4-diethenyl-3-methyl-, Copaene, (-)-1,2,2.alpha.,3,3,4,6,7,8,8.alpha.-decahydro-2.alpha.,7,8-trimethylacenaphthylene, beta.-Elemene, (Z)-.beta.-Caryophyllene, alpha.-ylangene, Humulene, beta.-copaene, Germacrene B, delta.-Cadinene, Elemol, Caryophyllene oxide, 5.alpha.,17.alpha.-Pregnan-12-one, Phytol, and Diisooctyl phthalate starting from a retention time of 6.6 min up to a retention time of 25.144 min. Beta-copaene was the major compound identified at a retention time of 12.91 min.

The compounds identified for *Hemigraphis alternata* are Undecane, 5,7-dimethyl-, 3-Ethyl-3-methylheptane, Cedrene-V6, Caryophyllene, Undecane, 3,8-dimethyl-, Hexane, 3,3-dimethyl-, Caryophyllene oxide, beta-Eudesmol, Diazoprogesterone, Diisooctyl phthalate. A total of 10 phytoconstituents was identified in the present study. Retention time taken for analysis was from 5.144 min to 25.179 min. The maximum height obtained was for peak 9 in the spectra representing diazoprogesterone at a retention time of 19.914 min. The prominent fractions identified in the present analysis are 3-Ethyl-3-methylheptane, Cedrene-V6, Caryophyllene, Caryophyllene oxide, Diazoprogesterone, Diisooctyl phthalate.



Chromatogram of *Chromolaena odorata*

Peak Report TIC							
Peak#	R.Time	Area	Area %	Height	Height %	Name	Base m/z
1	6.610	4881577	7.50	2709456	8.73	Cyclohexene, 5,6-diethenyl-1-methyl-	79.10
2	9.149	802073	1.23	394798	1.27	Cyclohexene, 3,4-diethenyl-3-methyl-	79.10
3	10.486	2744713	4.22	1433623	4.62	Copaene	105.20
4	10.613	417064	0.64	231917	0.75	(-)-1,2,2.alpha.,3,3,4,6,7,8,8.alpha.-decahydro-2.alpha.,7,8-trimethylacenaphthylene	189.20
5	10.715	927248	1.42	478471	1.54	.beta.-Elemene	93.15
6	11.232	10017124	15.39	5242829	16.90	(Z)-.beta.-Caryophyllene	93.10
7	11.373	376714	0.58	182378	0.59	.alpha.-ylangene	161.20
8	11.769	2213146	3.40	1152529	3.71	Humulene	93.15
9	12.191	27816313	42.74	14093841	45.42	.beta.-copaene	161.20
10	12.427	2433471	3.74	789017	2.54	Germacrene B	121.20
11	12.776	3999032	6.14	1932770	6.23	.delta.-Cadinene	161.20
12	13.196	1140621	1.75	459978	1.48	Elemol	59.05
13	13.785	903053	1.39	508291	1.64	Caryophyllene oxide	79.10
14	19.833	3359649	5.16	552403	1.78	5.alpha. 17.alpha. -Pregnan-12-one	233.15
15	20.396	1156278	1.78	336070	1.08	Phytol	71.10
16	25.144	1891730	2.91	529073	1.71	Diisooctyl phthalate	149.10
		65079806	100.00	31027444	100.00		


Component identification in *Chromolaena odorata*

Chromatogram of <i>Hemigraphis alternata</i> and component identification table							
Peak#	R.Time	Area	Area %	Height	Height %	Name	Base m/z
1	5.144	61094	2.15	37238	5.38	Undecane, 5,7-dimethyl-	57.10
2	8.721	139524	4.91	65000	9.39	3-Ethyl-3-methylheptane	57.10
3	10.622	136537	4.81	73174	10.57	Cedrene-V6	189.15
4	11.238	125156	4.41	61074	8.83	Caryophyllene	93.10
5	12.165	109270	3.85	48529	7.01	Undecane, 3,8-dimethyl-	57.10
6	12.849	91468	3.22	32186	4.65	Hexane, 3,3-dimethyl-	57.05
7	13.800	243185	8.56	98696	14.26	Caryophyllene oxide	79.15
8	14.815	176151	6.20	39854	5.76	.beta.-Eudesmol	59.15
9	19.914	1408141	49.58	158303	22.88	Diazprogesterone	233.10
10	25.179	349525	12.31	77906	11.26	Diisooctyl phthalate	149.10
		2840051	100.00	691960	100.00		


DISCUSSION

The GC-MS analysis conducted with the leaf samples of two plants revealed some compounds in common. The compounds identified in *Chromolaena odorata* are Cyclohexene, 5,6-diethenyl-1-methyl-, Cyclohexene, 3,4-diethenyl-3-methyl-, Copaene, (-)-1,2,2.alpha., 3,3,4,6,7,8.alpha.-decahydro-2.alpha., 7,8-trimethylacenaphthylene, beta.-Elemene, (Z)-.beta.-Caryophyllene, alpha.-ylangene, Humulene, beta.-copaene, Germacrene B, delta.-Cadinene, Elemol, Caryophyllene oxide, 5.alpha., 17.alpha.-Pregnan-12-one, Phytol, and Diisooctyl phthalate. The compounds identified for *Hemigraphis alternata* are Undecane, 5,7-dimethyl-, 3-Ethyl-3-methylheptane, Cedrene-V6, Caryophyllene, Undecane, 3,8-dimethyl-, Hexane, 3,3-dimethyl-, Caryophyllene oxide, beta.-Eudesmol, Diazprogesterone, Diisooctyl phthalate.


Reports earlier released,^[6-8] show the presence of monoterpenes, sesquiterpenes, hydrocarbons, triterpenes/steroids, alkaloids and flavonoids in *Eupatorium odorata*. Similarly, phenols, steroids and coumarins were detected in studies,^[9] conducted in *Hemigraphis alternata*. In concordance with the previous reports, both samples contain β -Caryophyllene/Caryophyllene, Caryophyllene oxide, Diisooctyl phthalate and class of steroid (Diazprogesterone/5- α , 17- α -Pregnan-12-one). Critical study of these compounds reveals the reason behind the wound healing property of the plants. β -caryophyllene, Caryophyllene and caryophyllene oxide belong to the sesquiterpene class of terpenoids.^[10]




β -caryophyllene^[11]



caryophyllene oxide^[11]



Diisooctyl phthalate^[12]



Diazprogesterone^[13]



5- α , 17- α -Pregnan-12-one^[14]

(Credit goes to online websites for molecular structures)

β -caryophyllene possess anti-inflammatory, antioxidant property. It is a bicyclic sesquiterpene, and according to studies can be used as alternative as aircraft fuel, therapeutic compound because of its anti-inflammatory and antioxidant nature.^[15] Diisooctyl phthalate belongs to phthalates (esters of o-phthalic acid) that has been detected in higher plants.^[16] Xenobiotic and pollutant aspects of phthalates were reported earlier.^[17] Work on the endogenous phthalates in plants has led to the view that phthalates has some physiological and biochemical role in plant protection.^[18] Diazprogesterone and 5- α , 17- α -Pregnan-12-one belongs to class of steroids. Steroids are reported as basic constituents of cell membranes they are said to function as signalling molecules in biological systems.^[19]

The antioxidant property shown by the terpenoids in both plants can be said to have significant function in wound healing process. Antioxidants reported in wound healing process,^[20] is in conformity with the finding. Fibroblast conservation and proliferation of keratinocyte in the wounded area due to the presence of antioxidants can be attributed to the wound healing efficiency of the plants.^[21] The phthalates, possess antimicrobial^[22] property, excretion of phthalates under stress by cells,^[23] ability of phthalates to reduce biofilm formation in bacterial phytopathogens^[24] all can be helpful in wound healing process exhibited in *Chromolaena odorata* and *Hemigraphis alternata*. The leaves of these plants may contain recommendable amount of phthalates which can be accounted for the reported results. The anti-inflammatory property of terpenoids is also a supporting action for wound healing. Infection, trauma, chemicals, heat, or unrecognized particles can induce inflammation of tissues.^[25] The influx of neutrophils and the resulting surge in free radicals can be alleviated by the antioxidant activity.^[26] Inflammatory disease pathology involves excessive Reactive Oxygen Species (ROS) generation by Polymorphonuclear Leukocytes (PMNLs). Natural compounds with antioxidant activity may be useful in reducing or regulating the oxidative damage caused by ROS produced from PMNLs. *C. odorata* may also contribute to wound healing by stimulating the production of antioxidants at the wound site, therefore protecting tissues from oxidative damage and providing a favourable environment for tissue healing.^[20] The same combination of terpenoid compounds detected in *Hemigraphis alternata* can be attributed to similar activity. Steroids are responsible for diverse physiological activities namely energy metabolism, circadian rhythms, inflammation, growth and reproduction, immunosup-

pression etc.,^[27] This property of steroids reported in the plants can be accounted for the blood clotting capability of the extracts. The infections coming in the wounded area may be prevented by the presence of phthalates as these compounds are playing major role in suppressing the growth and biofilm formation of bacterial phytopathogens.^[18] Based on the present study, the phytochemicals responsible for blood clotting and wound healing are identified. This study presents a preliminary comparative analysis of the two plants based on the phytochemicals and more precise quantitative determination of pharmacogenetic properties especially phthalates are to be conducted as these compounds are considered to be xenobiotic in scientific literature before utilization.

CONCLUSION

Present study aims to elucidate the common phytochemicals present in two plants *Chromolaena odorata* and *Hemigraphis alternata* that can be accounted for the wound healing properties. The presence of terpenes can be concluded as reason for the wound healing property as they have antioxidant and anti-inflammatory action; presence of steroids may be accounted for the blood clotting process and the presence of phthalates considered as preventing infection in the infected wound areas. Therefore, a proper method of isolation of these compounds and their optimal combination can help in instantaneous wound healing processes rather than moving on to the chemical synthesised combinations.

ETHICS APPROVAL

The submitted work is original and not placed for publication in any journal. It has not utilised any animal for study or pose threat to nature. Also, no external fund was utilised for the study and no additional contributors have been involved in the study.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ABBREVIATIONS

GC-MS: Gas Chromatography-Mass Spectrometry; **FTIR:** Fourier-Transform Infrared Spectroscopy; **ROS:** Reactive oxygen species; **PMNLs:** Polymorphonuclear leukocytes

SUMMARY

Two plants *Chromolaena odorata* and *Hemigraphis alternata* of family Asteraceae and Acanthaceae respectively were subjected to GC-MS analysis to analyse the common phytochemicals present that can be the responsible factor for the wound healing property of the plants that is commonly used in the local medicine in Kerala. The plant extracts of both were subjected to GC-MS analysis and the study revealed the presence of common components that can be accounted for the wound healing properties. The presence of terpenes can be concluded as reason for the wound healing property as they have antioxidant and anti-inflammatory action; presence of steroids may be accounted for the blood clotting process and the presence of phthalates considered as preventing infection in the infected wound areas. Therefore, a proper method of isolation of these compounds and their optimal combination can help in instantaneous wound healing processes rather than moving on to the chemical synthesised combinations.

REFERENCES

1. Ajay A, Rupesh MK, Shamal PAB, Abhishek K, Gowda, SK, Ramesh B. Pharmacological Importance of *Chromolaena odorata*: a review. Int J Pharm Drug Anal. 2021;9(1):8-11.
2. Yenti R, Afrianti R, Endang PA Formulasi Krim Ekstrak Etanol Daun Kirinyuh (*Eupatorium odoratum* L.) sebagai Antiinflamasi. Scientia: J Farmasi dan Kesehatan. 2016;4(1).
3. Abitha H, Prabhu K, Shakir A, George A, Aswathi P, Haritha P. A Study on Phytochemical Composition, GC-MS Analysis of ethyl acetate extract of *Hemigraphis alternata* IJCRT 2022;10(3):75-82.
4. Juszczak AM, Zovko-Končić M, Tomczyk M. Recent trends in the application of chromatographic techniques in the analysis of Luteolin and its derivatives. Biomolecules 2019;9:731.
5. Satapute P, Murali KP, Kurjogi M, Jogaiah S. Physiological adaptation and spectral annotation of Arsenic and Cadmium heavy metal-resistant and susceptible strain *Pseudomonas taiwanensis*. Environ. Pollut. 2019;251:555-63.
6. Akinmoladun AC, Ibukun EO, Don-Ologe IA. Phytochemicals constituents and antioxidant properties of extracts from the leaves of *Chromolaena odorata*. Sci Res Essay. 2007;2:191-4.
7. Zhang ML, Irwin D, Li XN, Sauriol F, Shi XW, Wang YF *et al.* PPAR γ agonist from *Chromolaena odorata*. J Nat Prod. 2012;5:2076-81.
8. Heiss EH, Tran TV, Zimmermann K, Schwaiger S, Vouk C, Mayerhofer B *et al.* Identification of chromomoric acid C-I as an Nrf2 activator in *Chromolaena odorata*. J Nat Prod. 2014;77:503-8.
9. Priya DM, Kumar VTG, Mathew F. Phytochemical screening of *Hemigraphis colorata* (Blume) H.G. Hallier J. Pharmacog. Phytochem. 2021;10(6):360-3.
10. Solomon H. The chemical and pharmacological basis of cloves (*Syzygium aromaticum* (L.) Merr. and L.M.Perry) as potential therapy for type 2 diabetes and associated diseases, In: Medicinal Foods as Potential Therapies for Type-2 Diabetes and Associated Diseases Editor(s): Solomon Habtemariam, Academic Press. 2019;551-78.
11. <https://www.sciencedirect.com/topics/pharmacology-toxicology-and-pharmaceutical->

12. <https://pubchem.ncbi.nlm.nih.gov/compound/Diisooctyl-phthalate#section=Structures>
13. https://pubchem.ncbi.nlm.nih.gov/compound/21-Diazopregn-4-ene-3_20-dione#section=Structures
14. https://pubchem.ncbi.nlm.nih.gov/compound/5alpha_17alpha-Pregnan-20-one_12beta-hydroxy#section=2D-Structure
15. Tundis R, Iacopetta D, Sinicropi MS, Bonesi M, Leporini M, Passalacqua NG., et al. Assessment of antioxidant, antitumor and pro-apoptotic effects of *Salvia fruticosa* Mill. subsp. *thomasi* (Lacaita) Brullo, Guglielmo, Pavone and Terrasi (Lamiaceae) *Food Chem Toxicol.* 2017;106:155-64.
16. Anantharaman A, Priya RR, Hemachandran H, Akella S, Rajasekaran C, Ganesh J., et al. Toxicity study of dibutyl phthalate of *Rubia cordifolia* fruits: *in vivo* and *in silico* analysis *Environmental toxicology* 2015;31.
17. Ventrice P, Ventrice D, Russo E, Sarro DG Phthalates: European regulation, chemistry, pharmacokinetic and related toxicity *Environ. Toxicol. Pharmacol.* 2013;36:88-96.
18. Shafikova TN, Maksimova LA, Omelichkina YV, Enikeev AG, Semenov AA. Endogenous phthalates in plants and their alleged participation in defense response against phytopathogens IOP Conference series: Earth and Environmental Science 2020;408:012076.
19. Sushma D, Parveen K, Gagandeep K. Biological Significance of Steroids, In: Steroids and their Medicinal Potential editor Abid Hussain Bandy; Bentham Science Publisher. 2023;98-124.
20. Cohen IK, Diegelmann RF, Lindblad WJ: Wound healing. *Biochem Clin Asp Philadelphia*; 1992.
21. Vaisakh MN, Pandey A. The invasive weed with healing properties: A review on *Chromolaena odorata*. *Int J Pharm Sci Res.* 2012;3:80-3.
22. Roy RN, Laskar S, Sen SK. Dibutyl phthalate, the bioactive compound produced by *Streptomyces albidoflavus* 321.2 *Microbiological research* 2006;161(2):121-6.
23. Babu B, Wu JT. Production of phthalate esters by nuisance freshwater algae and cyanobacteria. *Science of The Total Environment* 2010; 408(21):4969-75
24. Liu Y, Li J, Du J Accurate assessment of antibiotic susceptibility and screening resistant strains of a bacterial population by linear gradient plate *Sci China Life Sci* 2011;54(10):953-60.
25. Parimala DB, Tamilchelvan N, Ramasubramaniam R. Inflammation and medicinal plants-an ethnomedical approach. *J Phytol.* 2010;2:49-56.
26. Baboier BM. Oxygen-dependent microbial killing by phagocytes. *New Eng J Med.* 1978;29:629-68.
27. Crowley J, Withana M, Deplazes E. The interaction of steroids with phospholipid bilayers and membranes *Bio Phys Rev.* 2022;14(1):163-79.

Cite this article: Kundamveetil S. Comparative Phytochemical Analysis of Two Wound Healing Flora. *Asian J Biol Life Sci.* 2024;13(1):229-34.