

# Brine Shrimp Lethality and Antioxidant Activity of the Leaf, Rind and Seed Ethanolic Extracts of *Durio zibethinus* L.

Aileen May Gabule Ang\*, Cherry Mae Delos Reyes Nalda, Sandra Espinosa Sabejon

Department of Chemistry, College of Arts and Sciences, Central Mindanao University, Musuan, Bukidnon, PHILIPPINES.

Submission Date: 01-08-2018; Revision Date: 09-12-2018; Accepted Date: 14-12-2018

## ABSTRACT

**Introduction:** Substances derived from plants have recently receiving renewed interest due to their versatile applications. Parts of *Durio zibethinus* L. have been used traditionally as remedy to many diseases. To date, limited reports on the cytotoxicity and antioxidative property of *D. zibethinus* parts are available. **Objective:** The study aims to determine the cytotoxic and antioxidative property of the leaf, seed and rind ethanolic extracts of *D. zibethinus* L. **Materials and Methods:** Cytotoxicity of the leaf, seed and rind ethanolic extracts of *D. zibethinus* L. were determined through Brine Shrimp Lethality Assay (BSLA) while antioxidative property were measured via 2,2-diphenyl-1-picrylhydrazyl (DPPH) Radical Scavenging Activity and Reducing Power Assay. **Results and Discussions:** Results of the BSLA showed that the  $LC_{50}$  value of the ethanolic leaf, rind and seed extracts are 624.00, 139.31 and 46.28 mg/L, respectively. The data indicate potential cytotoxic property of the rind and seed extracts which can be attributed to the anthraquinones, flavonoids, phenolics, saponins and terpenoids detected in the rinds and seed extracts. The effective concentration ( $EC_{50}$ ) value for the DPPH radical scavenging activity and reducing power for the seed and rind ethanolic extracts showed significantly higher activity compared to the leaf extract. Similarities in the trend of the DPPH inhibition and reducing power of the ethanolic extracts of the leaves, rinds and seed ethanolic extracts imply direct correlation between DPPH scavenging activity and reducing power ( $P < 0.01$ ). **Conclusion:** *D. zibethinus* seeds and rinds can be potential sources of natural cytotoxic and antioxidant compounds.

**Key words:** *Durio zibethinus*, Cytotoxicity, Antioxidant property, DPPH, BSLA.

## Correspondence:

**Aileen May Gabule Ang**,  
Department of Chemistry,  
College of Arts and  
Sciences, Central  
Mindanao University, 8710  
Bukidnon, PHILIPPINES.  
Phone no: +639753344190.  
Email: aisemg\_ang@yahoo.  
com

## INTRODUCTION

Substances derived from plants have recently receiving renewed interest due to their versatile applications. Medicinal plants are abundant bioresource used in traditional and modern medications, nutraceuticals, food supplements, pharmaceutical intermediates and chemical entities for synthetic drugs.<sup>[1-2]</sup> Nowadays, researches are geared toward discovery of new drugs that possesses potency to combat the menace of drug resistant pathogenic microorganisms, antitumor and anticancer agents.

<sup>[3]</sup> Moreover, researches have indicated that the intake of plant-derived antioxidant helps prevent the development of some diseases. These antioxidants, which can be synthetic or natural, are vital to combat oxidative damage by free radicals in many oxidative stress-mediated disease conditions. However, synthetic antioxidants such as Butylated Hydroxyanisole (BHA) and Butylated Hydroxytoluene (BHT) have been recently reported to be dangerous to human health.<sup>[4]</sup>

Studies devoted to exploring plants as potential sources of natural cytotoxic antioxidant phytochemicals have been conducted. Phytochemicals or secondary metabolites, synthesized from plants, are capable of performing physiological actions in the body.<sup>[5-6]</sup> Some phytochemicals are found responsible to plants' antioxidant and cytotoxic properties.

## SCAN QR CODE TO VIEW ONLINE



www.ajbls.com

DOI :  
10.5530/ajbls.2018.7.13

*Durio zibethinus* L. is known as a king of fruits in South-east Asia with a distinctive thorn-covered husk and a unique odor. Of the known species that comprises the genus *Durio*, *D. zibethinus* L. is cultivated to a great extent.<sup>[7]</sup> About only 30% of the *D. zibethinus* fruit constitutes the only edible portion while the seeds and rinds are treated as food waste.<sup>[8]</sup> Moreover, numerous studies have reported that different parts of *D. zibethinus* L. trees have been traditionally used as an effective therapy against malaria fever, jaundice, skin diseases, swellings, colds, sores and wounds.<sup>[9]</sup> Different parts of *D. zibethinus* L. were also reported to exhibit anti-diabetic, antimicrobial, anti-inflammatory and anthelmintic properties.<sup>[10-13]</sup> However, limited reports on the cytotoxic and antioxidant activity among *D. zibethinus* L. plant parts are available. Thus, this study aimed to investigate the cytotoxic and antioxidative property of the leaf, seed and rind ethanolic extracts of *D. zibethinus* L. to provide scientific evidence that would support its medicinal properties.

## MATERIALS AND METHOD

### Sample Collection and Preparation

Mature leaves and fruits of *Durio zibethinus* L. were collected from Patag, Alanib, Lantapan, Bukidnon with Global Positioning System (GPS) coordinates of 8°01'57.9" N and 124°58'56.3" E. The leaf samples were washed with distilled water to remove dirt. The rinds and seeds of the fruit were separated from the flesh and washed with distilled water. Identification and authentication of the samples were done at the Botany section of the Central Mindanao University (CMU) Museum, Musuan, Bukidnon. The samples were then air-dried separately until the moisture content is reduced to at least 10 percent. The dried samples were pulverized using a grinder, extracted with ethanol for 72 hrs and filtered. The filtrates were then concentrated in vacuo at 40°C to yield Concentrated Leaf (LE), Rind (RE) and Seed (SE) ethanolic extracts.

### Phytochemical Screening

The LE, RE and SE were screened for presence of phytochemicals, i.e. alkaloids, anthraquinones, flavonoids, phenols, saponins, tannins and terpenoids employing Thin-layer Chromatography (TLC) method with slight modification.<sup>[14]</sup>

### Brine Shrimp Lethality Assay

The brine shrimp lethality assay was carried out with slight modification.<sup>[15]</sup> A 10,000 mg/L LE, RE and SE stock solutions in ethanol were prepared. Appropriate

volumes of 500.0, 250.0, 50.0, 25.0, 5.0 and 2.5 µL of the stock solution were placed into separate test tubes marked up to 5.00 mL to give 1000, 500, 100, 50, 10 and 5 mg/L, respectively. The solution was purged with N<sub>2</sub> gas to remove the solvent. Minimal amount of Dimethyl Sulfoxide (DMSO) corresponding to 50.00, 25.00, 5.00, 2.50, 0.50 and 0.25 µL for 1000, 500, 100, 50, 10 and 5 mg/L, respectively, were added as liquid surfactant into each test tube. The mixture was then vortexed to ensure homogeneity. Ten brine shrimp nauplii were transferred from the hatching chamber into each test tubes. Sterilized seawater was added up to the 5 mL mark and the test tubes were incubated under illumination for 24 hrs. Negative and positive controls were run using sterilized seawater with DMSO and potassium dichromate, respectively. The number of surviving shrimps were counted and recorded after 24 hrs. The percentage mortality of nauplii was then calculated.

For sample test solutions giving 0% and 100% mortalities, necessary corrections were done.<sup>[16]</sup> Using the Probit Table of Finney, the corrected percentage mortality values were transformed to probit values.<sup>[17]</sup> The best-fitting straight curve was then drawn with the logarithm of the concentration values as the abscissa and the probit values as the ordinate. The log<sub>10</sub> concentration value that corresponds to the probit point of 5.00 from the curve was recorded and converted to its antilog value to give the LC<sub>50</sub>. LC<sub>50</sub> of less than 200 mg/L was considered cytotoxic.<sup>[18]</sup> Analysis was carried out in five replicates. Result was then reported as mean of the analysis.

### DPPH Radical Scavenging Activity Assay

The DPPH radical scavenging activity assay was carried out in a 96-well microplate.<sup>[19]</sup> Various concentrations of LE, RE and SE test solutions (5 -500 µg/mL) were prepared in ethanol.

In a well containing 150 µL of sample solution, 50 µL of 0.1 mM DPPH in absolute ethanol was added. The mixture was incubated for 30 mins in the dark at room temperature. The absorbance for the reaction mixtures was measured at 517 nm using a microplate reader. The same procedure was done for the various gallic acid solutions (15-210 µg/mL) and absolute ethanol (blank). The absorbance readings were recorded and the DPPH radical scavenging activity were calculated as percentage DPPH inhibition (Equation 1).

$$\% \text{ DPPH inhibition} = \left( \frac{A_{\text{cont}} - A_{\text{sam or std}}}{A_{\text{cont}}} \times 100 \right) \quad \text{Equation 1}$$

where:

A<sub>control</sub> = Absorbance of the control

A<sub>sample</sub> = Absorbance of the sample solution

Astandard = Absorbance of the gallic acid solution

The Least Squares Method was then employed to obtain the equation of the line. The equation was used to calculate the EC<sub>50</sub> (concentration required to obtain a 50% DPPH radical inhibition) of the extract solutions. Result is the mean value of the triplicate analysis.

### Reducing Power Assay

The reducing power was determined employing the method adapted for a 96-well plate assay.<sup>[20]</sup> Various concentrations of SE, RE and SE test solutions (50, 100, 250, 500 and 750 µg/mL) were prepared in ethanol. In an Eppendorf tube containing 1 mL of the sample test solution, 200 µL of 0.2 M phosphate buffer (pH 6.6) and 200 µL of 1% (w/v) solution of potassium ferricyanide were added. The mixture was then incubated at 50°C for 30 mins. Then 10% (w/v) trichloroacetic acid (200 µL) was added after the solution was cooled to room temperature. The resulting mixture was centrifuged at 11000 rpm for 3 mins. After centrifugation, an aliquot (200 µL) of the upper layer of the solution was transferred to a 96-well plate and 20 µL of 0.1% (w/v) solution of ferric chloride was added. The absorbance of the solution was measured at 620 nm using a microplate spectrophotometer reader. The same procedure was done for the negative control (absolute ethanol) and standard gallic acid test solutions (15, 30, 45, 60, 75 and 90 µg/mL). The percent reducing power of the sample and gallic acid test solutions were then calculated using Equation 2.

$$\text{Reducing Power (\%)} = \left( \frac{A_{\text{sample or std}} - A_{\text{cont}}}{A_{\text{sample}}} \times 100 \right) \quad \text{Equation 2}$$

where:

Acontrol = Absorbance of the control

Asample = Absorbance of the sample test solution

The EC<sub>50</sub> (mg/L) of the sample solutions was determined by Least Squares Method. The equation of the line was then used to calculate the EC<sub>50</sub> (concentration required to obtain a 50% DPPH radical inhibition) of the solutions. Result is the mean value of the triplicate analysis.

## RESULTS

### Phytochemical Screening

The results of the phytochemical screening are shown in Table 1.

### Brine Shrimp Lethality

The percent mortality of the *A. salina* nauplii in LE, RE and SE and the LC<sub>50</sub> of the sample extracts are presented in Table 2.

**Table 1: Phytochemical screening of ethanolic leaf, rinds and seed extracts of *D. zibethinus* L.**

PHYTOCHEMICAL	ETHANOLIC EXTRACT			Flesh <sup>[21]</sup>
	Leaves	Rinds	Seeds	
Alkaloids	+	-	-	+
Anthraquinones	-	+	+	+
Flavonoids	-	+	+	+
Phenolics	+	+	+	+
Saponins	-	+	+	+
Tannins	-	-	-	+
Terpenoids	-	+	+	+


(+) – present; (-) – absent.

**Table 2: LC<sub>50</sub> of the extracts and percent mortality (%) of the brine shrimp nauplii after 24hr exposure to the ethanolic leaf, rind and seed extracts of *D. zibethinus* L.**


Sample	Mortality* (%)						LC <sub>50</sub> (mg/L)	Inference**
	5 mg/L	10 mg/L	50 mg/L	100 mg/L	500 mg/L	1000 mg/L		
Leaves	2.50 ± 0.00	2.50 ± 0.00	2.50 ± 0.00	19.50 ± 2.09	43.00 ± 10.06	63.00 ± 2.74	624.00	Non-cytotoxic
Rinds	3.25 ± 0.00	11.88 ± 2.46	30.00 ± 1.77	46.00 ± 9.45	70.63 ± 2.39	85.38 ± 4.89	139.31	Cytotoxic
Seeds	10.32 ± 0.36	25.50 ± 3.26	50.00 ± 10.16	68.50 ± 4.19	83.63 ± 7.20	95.50 ± 2.87	46.28	Cytotoxic
Positive Control (K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> )	90.00 ± 0.00	97.50 ± 0.00	97.50 ± 0.00	97.50 ± 0.00	97.50 ± 0.00	97.50 ± 0.00	< 5.00	Cytotoxic

\*Data are expressed as mean ± SD. Each sample was analyzed five times.

\*\*LC<sub>50</sub> < 200 mg/L is cytotoxic.[18]



**Figure 1:** EC<sub>50</sub> value (mg/L) for the DPPH radical scavenging activity of the ethanolic extracts of *D. zibethinus* L. leaves, rinds and seeds. Error bars are standard error (n=3).



**Figure 2:** EC<sub>50</sub> value (mg/L) for the reducing power of the ethanolic extracts of *D. zibethinus* L. leaves, rinds and seeds. Error bars are standard error (n=3).

### DPPH Radical Scavenging Activity

The EC<sub>50</sub> for the DPPH radical scavenging activity of the *D. zibethinus* ethanolic extracts are shown in Figure 1. The mean values of the same letter superscript are not significantly different by Tukey's Test at 0.05 level of significance.

### Reducing Power

The antioxidant property of *D. zibethinus* in terms of reducing power is presented in Figure 2. The mean values of the same letter superscript are not significantly different by Tukey's Test at 0.05 level of significance.

## DISCUSSION

### Phytochemical Screening

Alkaloids and phenolics were detected in the ethanolic leaf extract of *D. zibethinus* L., while anthraquinones,

flavonoids, phenolics, saponins and terpenoids were detected in the ethanolic rind and seed extracts (Table 1). The positive results of the screening are consistent with the results of the previous studies using the flesh of *D. zibethinus* L.<sup>[12,21]</sup>

Several studies have reported that alkaloids, anthraquinones, flavonoids, phenolics, saponins and terpenoids possess biological properties such as antiapoptosis, antiaging, anticarcinogen, antiinflammation, antiatherosclerosis and cardiovascular protection<sup>[3,22,23]</sup> as well as analgesic, antispasmodic, antibacterial, antimutagenic, antioxidant and antiviral effects.<sup>[13,24-25]</sup> These findings may support claims on the reported medicinal properties of the leaves, rinds and seeds of *D. zibethinus* L. such as lipid lowering effect, immunomodulatory and antibacterial effect, efficiency in interaction with medicinal entities, bactericidal and antibacterial effects, antimicrobial and antiviral activities, synergism effect with Penicillin G and antioxidative activities.<sup>[26-29]</sup>

### Brine Shrimp Lethality

The Brine Shrimp Lethality Assay (BSLA) was carried out to evaluate the cytotoxicity of the ethanolic leaf, rind and seed extracts of *D. zibethinus* L. against the brine shrimp nauplii. The result of BSLA is summarized in Table 2.

The percent mortality of the brine shrimp nauplii increases as the concentration of the test solutions are increased. Apparently, the percent mortality of the nauplii was in the order of leaves < rinds < seeds extracts for all concentrations of the test solutions. This indicates that the mortality of the nauplii in sample extracts is concentration-dependent. The highest percent mortality values in 5 mg/L to 1000 mg/L test solutions were consistently recorded for the ethanolic seed extracts while the lowest percent mortality values were observed for the ethanolic leaf extracts. These findings indicate that the ethanolic extracts of the various plant parts

**Table 3: Pearson's correlation test between DPPH radical scavenging activity and reducing power of the ethanolic extracts of *D. zibethinus* L. leaves, rinds and seeds.**

		DPPH	Reducing Power
DPPH	Pearson Correlation	1	.950**
	Sig. (2-tailed)		<.001
	N	9	9
Reducing Power	Pearson Correlation	.950**	1
	Sig. (2-tailed)	<.001	
	N	9	9

\*\* Correlation is significant at the 0.01 level (2-tailed).



of *D. zibethinus* L. exhibit cytotoxicity against the brine shrimp nauplii and that its toxicity to the nauplii is plant part-dependent. The relatively high cytotoxicity of the ethanolic rind and seed extracts than the leaf extracts can be due to the different phytochemicals in *D. zibethinus* L. rinds and seeds. Apart from the alkaloids and phenolics which are present among the three plant parts, the seeds and the rind ethanolic extracts contain anthraquinones, flavonoids, saponins and terpenoids as well. Studies have shown that these phytochemicals are known to exhibit analgesic, antispasmodic, antibacterial, antimicrobial, anti-inflammatory, antioxidant, anti-allergenic, diuretic, antiviral, anticancer and antitumor properties.<sup>[14,30-33]</sup>

For the  $LC_{50}$ , the ethanolic leaf extract gave the highest value (624.00 mg/L), followed by the rind (139.31 mg/L) and the seed (46.28 mg/L). Since the  $LC_{50}$  value and the plant extract toxicity are inversely proportional, the cytotoxic property of the plant parts increases in order of leaves < rinds < seeds. Plant extracts with  $LC_{50}$  less than 200 ppm are considered significantly active and had the potential for further investigation.<sup>[18]</sup> Thus, rinds and seeds exhibited potential cytotoxicity that would warrant further studies.

### DPPH Radical Scavenging Activity

The results of the determination of the DPPH radical scavenging activity, in terms of  $EC_{50}$  value, of the ethanolic extracts of *D. zibethinus* L. leaves, rinds and seeds show that among the plant parts, the leaves gave the highest  $EC_{50}$  value, followed by the rinds and seeds which recorded of 229.29, 54.14 and 44.17 mg/L respectively (Figure 1). On the other hand, pure gallic acid recorded a value of less than 15.00 mg/L. The result show that *D. zibethinus* L. leaves, rinds and seeds exhibit antioxidative property via DPPH free radical scavenging mechanism.

The  $EC_{50}$  is the efficient concentration of the substrate which is required to cause an amount of 50% loss in DPPH radical activity.<sup>[34]</sup> The lower the  $EC_{50}$  value, the higher the scavenging activity.<sup>[35]</sup>

The results of the Analysis of Variance (ANOVA) at 0.05 level of significance and the subsequent Post Hoc Tukey's Test indicate that the ethanolic extracts of *D. zibethinus* L. rinds and seeds have statistically similar  $EC_{50}$  value but is significantly lower than that of *D. zibethinus* L. leaves. This indicate that the rind and seed extracts have higher DPPH radical scavenging activity than the leaf extract. Thus, the DPPH radical scavenging activity of *D. zibethinus* L. is plant-part dependent.

Phytochemicals in different plant parts is speculated to affect their bioactivity. This was demonstrated in a

study wherein total phenolic and flavonoid content and antioxidant activity of *Betula utilis* D. Don vary among plant parts.<sup>[36]</sup> Moreover, these findings support results of previous studies which detected presence of polyphenols like phenolics and flavonoids in the pulp of *D. zibethinus* which are contributors to the plant's antioxidant capacity.<sup>[37]</sup> Presence of polyphenol and Superoxide dismutase (SOD) in the durian seed, sarcocarp and peel extracts.<sup>[37-38]</sup>

### Reducing Power

Reductive capabilities of plant extracts can serve as a significant indicator of their potential antioxidant activities.<sup>[39]</sup>

As shown in Figure 2, seed ethanolic extract exhibited the highest reducing power followed by rind and leaf extracts. However, pure gallic acid showed higher activity. Results imply that *D. zibethinus* L. leaves, rinds and seeds possess antioxidative property specifically reduction capability.

The results of the Analysis of Variance (ANOVA) at 0.05 level of significance and the subsequent Post Hoc Tukey's Test reveal that *D. zibethinus* L. rind and seed ethanolic extracts demonstrated significantly higher reducing power and consequently higher antioxidant activity. This can be accounted to the differences in the type and amount or concentration of phytochemicals present in the different plant parts.<sup>[40]</sup> Thus, antioxidant activity, of *D. zibethinus* L., in terms of reducing power, vary with plant part.

However, the similarities in the trend of the DPPH inhibition and reducing power in *D. zibethinus* L. leaves, rinds and seeds is interesting to note. Using Pearson's correlation test, which quantifies the linear association between two quantitative variables, it is possible to estimate the relationship between DPPH inhibition and reducing power. The results, as shown in Table 3, reveal that there is a positive correlation between DPPH and reducing power ( $P < 0.01$ ) in *D. zibethinus* L. leaves, rinds and seeds. This implies that increase in reducing power results in a corresponding increase in the DPPH radical scavenging activity of the ethanolic extracts of *D. zibethinus* L. leaves, rinds and seeds.

The same correlation between DPPH radical scavenging activity and reducing power was observed in *Porphyra indica* ( $P < 0.01$ ) and *Porphyra vietnamensis* ( $P < 0.05$ ).<sup>[41]</sup> This observation indicates that the substances responsible to reduce  $Fe^{3+}$  to  $Fe^{2+}$  are capable of scavenging DPPH radical satisfactorily and suggest that these substances might react with DPPH radical through electron transfer reaction mechanism.<sup>[42]</sup>

## CONCLUSION

The present data obtained showed that ethanolic rind and seed extracts of *D. zibethinus* possess cytotoxic and antioxidant properties. Results of the study may also serve as scientific evidence to support traditional use of *D. zibethinus* L. rinds and seeds. Rinds and seeds of *D. zibethinus* L., which are considered as food waste, may be utilized as potential source of natural cytotoxic and antioxidant compounds. Thus, studies to further establish the bioactivity of *D. zibethinus* L. and to isolate the compounds responsible for its bioactivity is recommended.

## ACKNOWLEDGMENT

The authors wish to thank the Chemistry Department, Natural Science Research Center and Natural Products Research and Development Center of Central Mindanao University, Musuan, Bukidnon for the assistance during the conduct of the study.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

## ABBREVIATIONS

**ANOVA:** Analysis of Variance; **BHA:** Butylated; hydroxyanisole; **BHT:** Butylated hydroxytoluene; **BSLA:** Brine Shrimp Lethality Assay; **DPPH:** 2,2-diphenyl-1-picrylhydrazyl; **DMSO:** Dimethylsulfoxide; **EC<sub>50</sub>:** Effective Concentration; **LC<sub>50</sub>:** Lethal Concentration; **SOD:** Superoxide dismutase; **TLC:** Thin Layer Chromatography.

## SUMMARY

The study evaluates the cytotoxic and antioxidative property of the leaf, seed, and rind ethanolic extracts of *Durio zibethinus* L. The results obtained indicate potential cytotoxic property of the rind and seed ethanolic extracts. Moreover, a significantly higher DPPH radical scavenging activity and reducing power was exhibited by the seed and rind extracts as compared to the leaf extracts. These can be attributed to the phytochemicals, i.e. anthraquinones, flavonoids, phenolics, saponins, and terpenoids, detected in the said plant extracts. Thus, *D. zibethinus* seeds and rinds can be potential sources of natural cytotoxic and antioxidant compounds.

## REFERENCES

- Pandey A, Tripathi S. Concept of standardization, extraction and pre phytochemical screening strategies for herbal drug. J Pharmacogn Phytochem. 2014;2(5):115-9.
- Tiwari P, Kumar B, Kaur M, Kaur G, Kaur H. Phytochemical screening and extraction: a review. Int Pharm Sci. 2011;1(1):98-106.
- Musa AA. Cytotoxicity activity and phytochemical screening of *Cochlospermum tinctorium* perr ex A. Rich rhizome. J Appl Pharm Sci. 2012;2(7):155-9.
- Gupta VK, Sharma SK. Plants as natural antioxidants. Nat Prod. 2006;5(4):326-34.
- Kumar BR. Application of HPLC and ESI-MS techniques in the analysis of phenolic acids and flavonoids from Green Leafy Vegetables (GLVs). J Pharm Anal. 2017;7(6):349-64.
- Iqbal E, Salim KA, Lim LBL. Phytochemical screening, total phenolics and antioxidant activities of bark and leaf extracts of *Goniothalamus velutinus* (Airy Shaw) from Brunei Darussalam. J King Saud Univ. 2015;27(3):224-32.
- Lim TK. Durian Diseases and Disorders. Tropical Press, SDN. BHD. Kuala Lumpur. 1990;107.
- Amin AM, Ahmad AS, Yin YY, Yahya N, Ibrahim N. Extraction, purification and characterization of durian (*Durio zibethinus*) seed gum. Food Hydrocolloids. 2007;21(2):273-9.
- Brown MJ. Durio - A bibliographic review. New Delhi: International Plant Genetic Resources Institute (IPGRI) office for South Asia. 1997.
- Chingsuwanrote P, Muangnoi C, Parengam K, Tuntipopipat S. Antioxidant and anti-inflammatory activities of durian and rambutan pulp extract. Int Food Res J. 2016;23(3):939-47.
- Primarianti AU, Sujono TA. Antidiabetic activity of durian (*Durio zibethinus* Murr.) and rambutan (*Nephelium lappaceum* L.) fruit peels in alloxan diabetic rats. Procedia Food Sci. 2015;3:255-61.
- Sah BP, Pathak T, Sankar S, Suresh B. Phytochemical investigations on the fruits of *Durio zibethinus* Linn. for antimicrobial activity. Int J Pharma Sci Res. 2014;5(12):878-91.
- Ashraf MA, Maah MJ, Yusoff I. Estimation of antioxidant phytochemicals in four different varieties of durian (*Durio zibethinus* Murray) fruit. Middle East J Sci Res. 2010;6(5):465-71.
- Aguinaldo AM, Espeso EI, Guevarra BQ, Nonato MG. A guidebook to plant screening: Phytochemical and biology. Phytochemistry section. Manila: UST Publishing House. 2005.
- McLaughlin JL. Crown gall tumours on potato discs and brine shrimp lethality: two simple bioassays for higher plant screening and fractionation. Methods in Plant Biochemistry. 1991;6:1-32.
- Vincent K. Probit analysis. San Francisco: San Francisco State University. 2008. In: <http://userwww.sfsu.edu/efc/courses/biol710/probit/Probit-Analysis.htm>.
- Finney D. Probit analysis: A statistical treatment of the sigmoid response curve. Cambridge University Press, Cambridge. 1952.
- Alali FQ, Tawaha K, El-Elmat T, Qasaymeh R, Li C, Burgess J, et al. Phytochemical studies and cytotoxicity evaluations of *Colchicum tunicatum* Feinbr and *Colchicum hierosolymitanum* Feinbr (Colchicaceae): two native Jordanian meadow saffrons. Natural Product Research. 2006;20(6):558-66.
- Miser-Salihoglu E, Akaydin G, Caliskan-Can E, Yardim-Akaydin S. Evaluation of antioxidant activity of various herbal folk medicines. J Nutr and Food Sci. 2013;3:222-31.
- Murugan K, Iyer VV. Antioxidant and antiproliferative activities of Marine Algae, *Gracilaria edulis* and *Enteromorpha lincolata*, from Chennai Coast. Int J Cancer Res. 2012;8(1):15-26.
- Ho LH, Bhat R. Exploring the potential nutraceutical values of durian (*Durio zibethinus* L.)—An exotic tropical fruit. Food Chem. 2015;168:80-9.
- Aiyegoro OA, Okoh AI. Preliminary phytochemical screening and *in vitro* antioxidant activities of the aqueous extract of *Helichrysum longifolium* DC. BMC Compl and Alt Med. 2010;10(1):21.
- Motaung SCKM, Razwinani M, Tshikalange ET. Phytochemical screening and cytotoxicity activity of crude extracts from *Pleurostylia capensis* (celastraceae). Biochem and Pharmacol. 2013;2(4):79.
- Sülse VP, Lizarraga E, Mamadaliyeva NZ, Lago JH. Potential of Terpenoids and Flavonoids from Asteraceae as Anti-Inflammatory, Antitumor and Antiparasitic Agents. Evidence-Based Complementary and Alternative Medicine. 2017;2017:1-2.
- Mohammed RM, Ayoub SM. Study of Phytochemical Screening and Antimicrobial Activity of Citrus aurantifolia Seed Extracts. Am J Analyt Chem. 2016;7(3):254-9.

26. Manoharan S. Synergistic activity of chloroform extract of *Durio zibethinus* wood bark with penicillins against *Staphylococcus aureus*. Int J Biol Med Res. 2013;4(2):3025-7.
27. Pholdaeng K, Pongsamart S. Studies on the immunomodulatory effect of polysaccharide gel extracted from *Durio zibethinus* in *Penaeus monodon* shrimp against *Vibrio harveyi* and WSSV. Fish and Shellfish Immunol. 2010;28(4):555-61.
28. Toledo F, Arancibia-Avila P, Park YS, Jung ST, Kang SG, GuHeo B, et al. Screening of the antioxidant and nutritional properties, phenolic contents and proteins of five durian cultivars. Int J Food Sci Nutr. 2008;59(5):415-27.
29. Voon YY. Physicochemical Properties of Whole and Fresh Cut Durian (*Durio Zibethinus*) Stored at Ambient and Refrigerated Temperatures (Doctoral dissertation, Universiti Putra Malaysia). 2008.
30. Aziz MA, Alam AS, Ema AA, Akter M, Chowdhury MM. Analysis of secondary metabolites, antibacterial, brine shrimp lethality and larvicidal potentiality of *Microcos paniculata* fruits. IOSR J Pharm Biol Sci. 2014;9(3):50-8.
31. Odeja OO, Obi G, Ogwuche CE, Elemike EE, Oderinlo OO. Phytochemical screening, Antioxidant and Antimicrobial activities of *Senna occidentalis* (L.) leaves. Int J Herb Med. 2014;2(4):26-30.
32. Sharma G, Srivastava AK, Prakash D. Phytochemicals of Nutraceutical importance: their role in health and diseases. Pharmacologyonline. 2011;2:408-27.
33. Thoppil RJ, Bishayee A. Terpenoids as potential chemopreventive and therapeutic agents in liver cancer. World J Hepatol. 2011;3(9):228-49.
34. Molyneux P. The use of the stable free radical Diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. Songklanakarin J Sci Technol. 2004;26(2):211-9.
35. Omenna CE. Antioxidative activity of the almond leaves (*Terminalia catappa*). Int J Nurs Midwifery. 2015;1(2):29-40.
36. Wani MS, Gupta RC, Munshi AH, Pradhan SK. Phytochemical screening, total phenolics, flavonoid content and antioxidant potential of different parts of *Betula utilis* Don from Kashmir Himalaya. Int J Pharm Sci Res. 2018;9(6):2411-7.
37. Ji YS, Chang JP. Antioxidative activity of the durian (*Durio zibethinus*) extract. Korean J Med Crop Sci. 2013;21(4):255-61.
38. Duazo NO, Bautista JR, Teves FG. Crude methanolic extract activity from rinds and seeds of native durian (*Durio zibethinus*) against *Escherichia coli* and *Staphylococcus aureus*. Afr J Microbiol Res. 2012;6(35):6483-6.
39. DeAncos B, González EM, Cano MP. Ellagic acid, vitamin C and total phenolic contents and radical scavenging capacity affected by freezing and frozen storage in raspberry fruit. J Agric Food Chem. 2000;48(10):4565-70.
40. Raja R, Hemaiswarya S, Arunkumar K, Carvalho IS. Antioxidant activity and lipid profile of three seaweeds of Faro, Portugal. Braz J Bot. 2016;39(1):9-17.
41. Pise NM, Jena KB, Maharana D, Gaikwad D, Jagtap TG. Free radical scavenging potential, reducing power, phenolic and biochemical constituents of *Porphyra* species from India. J Algal Biomass Utiln. 2010;1(3):29-42.
42. Košmerl T, Bertalančič L, Maras V, Kodžulović V, Šučur S, Abramovič H. Impact of yield on total polyphenols, anthocyanins, reducing sugars and antioxidant potential in white and red wines produced from Montenegrin autochthonous grape varieties. J Food Sci Technol. 2013;1(1):7-15.

**Cite this article:** Ang AMG, Nalda CMDR, Sabeyon SE. Brine Shrimp Lethality and Antioxidant Activity of the Leaf, Rind and Seed Ethanolic Extracts of *Durio zibethinus* L. Asian J Biol Life Sci. 2018;7(3):105-11.