

A Review on Recognition of Plant Disease using Intelligent Image Retrieval Techniques

Gulbir Singh^{1,2,*}, Kuldeep Kumar Yogi¹

¹Department of Computer Science, Banasthali Vidyapith, Rajasthan, INDIA.

²MMICT & BM, M.M. (Deemed to be University), Ambala, Haryana, INDIA.

Submission Date: 04-11-2020; Revision Date: 30-11-2020; Accepted Date: 10-12-2020

ABSTRACT

Today, crops face many characteristics/diseases. Insect damage is one of the main characteristics/diseases. Insecticides are not always effective because they can be toxic to some birds. A common practice of plant scientists is to visually assess plant damage (leaves, stems) due to disease based on the percentage of disease. Plants suffer from various diseases at any stage of their development. It requires urgent diagnosis and preventive measures to maintain quality and minimize losses. Many researchers have provided plant disease detection techniques to support rapid disease diagnosis. In this review paper, we mainly focus on artificial intelligence (AI) technology, image processing technology (IP), deep learning technology (DL), vector machine (SVM) technology, the network Convergent neuronal (CNN) content detailed description of the identification of different types of diseases in tomato and potato plants based on image retrieval technology (CBIR). It also includes the various types of diseases that typically exist in tomatoes and potatoes. Content-based Image Retrieval (CBIR) technologies should be used as a supplementary tool to enhance search accuracy by encouraging you to access collections of extra knowledge so that it can be useful. CBIR systems mainly use colour, form and texture as core features, such that they work on the first level of the lowest level. This is the most sophisticated method used to diagnose diseases of tomato plants.

Key words: Artificial Intelligence, Deep Learning, Tomato Leaf Disease, Potato Leaf Disease, Content-based Image Retrieval (CBIR).

Correspondence:

Mr. Gulbir Singh^{1,2}

¹Scholar, Department of Computer Science, Banasthali Vidyapith, (Rajasthan), INDIA.

²MMICT & BM, M.M. (Deemed to be University), Ambala, (Haryana), INDIA.

Phone no: +91-09719042509

Email: gulbir.rkgit@gmail.com

INTRODUCTION

Agriculture is the backbone of the economy of India. The large-scale commercialisation of agriculture has had a very negative effect on our climate. The use of chemical pesticides creates a substantial build-up of chemicals in our atmosphere, soil, water, food, livestock and also in our own bodies. Artificial fertilisers have a short-term effect on production but have a long-term detrimental impact on the environment. After leaching and loss, the chemical fertiliser will remain in the soil for many years and will pollute the groundwater. Another

negative effect of this development is its effect on the situation of rural populations around the world. Despite the so-called rise in production, the plight of farmers in nearly every country in the world is deteriorating. It's the root of organic farming. Natural farming can solve all these issues. The main practices in organic farming rely on fertilisation and control of pests.^[1]

Potato is one of the most significant food crops. The diseases causing substantial yield loss in potato are *Phytophthora infestans* (late blight) and *Alternaria solani* (early blight). Early detection of these diseases can allow to take preventive measures and mitigate economic and production losses. Over the last decades, the most practiced approach for the detection and identification of plant diseases is naked eye observation by experts. But in many cases, this approach proves unfeasible due to the excessive processing time and unavailability of experts at farms located in the remote areas. Hence,

SCAN QR CODE TO VIEW ONLINE



www.ajbls.com

DOI: 10.5530/ajbls.2020.9.42

the introduction of image analysis tools turns out to be an effective method for continuous monitoring of plant health status and early detection of plant diseases. As diseases leave some visible symptoms on the plants, particularly on leaves, disease detection can be performed by imaging analysis of those visible patterns on leaves. Thus imaging technique combined with machine learning offers a solution to the issue of agricultural productivity and ensures food security.^[2]

Detection of plant diseases by physically analysing the signs of plant leaves easily brings difficulty. Due to this difficulty and the vast number of crops grown and plant pathology issues that occur, even qualified agricultural specialists and plant pathologists will sometimes struggle to identify particular diseases, leading to inaccurate findings and responses to program issues. An automated device designed to help diagnose plant diseases by the presence and visual signs of plants can be of great benefit to agricultural production enthusiasts. This will prove to be a valuable strategy for farmers and will be recalled at the right time before the disease spreads over a wide region.^[3]

Potato is one of the most important crops for food. The diseases that have induced a sharp decline in potato production are *Phytophthora* (downy mildew) and *Streptomyces solani* (downy mildew). Early identification of these diseases can take preventive measures to minimise economic and production losses. Over the past decades, visual evaluation by professionals has been the most realistic way of detecting and distinguishing plant diseases. In certain cases, however, this approach has proved to be inefficient due to the long processing period and inaccessibility of agricultural experts in remote areas. The implementation of image processing software has also proved to be an important approach for the continuous monitoring of plant health and early identification of plant diseases. If the disease leaves noticeable signs on the plant, particularly on the leaves, the disease can be identified by imaging and examining these recognisable patterns on the leaves. Imaging technology combined with machine learning also offers a solution to the issue of agricultural production and assures food security.

Tomato is one of the world's most valuable crops. According to the Food and Agriculture Organisation of the United Nations, the production of world tomatoes amounted to approximately 170,750 kilotons in 2014. Plant diseases have always been a thorny problem in agricultural production and one of the key factors restricting sustainable agricultural growth. Tomato is a popular vegetable and a major economic crop in China. It is extensively planted in different areas and

occupies an area of around 700 million square meters. Tomato disease is also caused by different environmental factors.^[4] Detection of plant diseases is always a daunting task. The major sources of infection are fungi, bacteria and viruses in plants. These infections can affect any part of the plant, such as leaves, stems and roots.

In order to obtain better quality and profitable crops, farmers recognise the right goods by measuring and regulating the required specifications for temperature, light and humidity.^[5] In addition, due to population growth, climate change and political instability, agriculture has begun to find new ways to expand food production. This encourages researchers to look for new, resource-rich and efficient innovations that can help improve agricultural productivity. However, difficulties exist, such as early identification of pathogens in plants and plant problems. It is difficult to continually detect the kinds of diseases in the leaves of plants with the naked eye. The electronic system of specialists would also be very helpful and will aid in the timely diagnosis of diseases. Alternative electronic approaches for quick and non-invasive detection of tomato diseases have been investigated in recent years.

Various diseases generally occur in plants Figure 1;

Precision cultivation can be used to control certain pests and diseases. The use of these chemical or expensive processes can be minimised by using precise cultures. In the field of precision agriculture, sensor technology, computer processing, remote sensing processing,




Figure 1: Various causes of disease in plants.

robotics and other information technologies are used. For precision agricultural applications^[6] (such as spraying only infected areas), it is important to recognise places where plant diseases are occurring and spread. Operators, static stations, sensor networks, drones and mobile robots are used for the monitoring of precision agriculture. The only drawback of these methods is that they cannot find specialists in the area. In order to conduct precision breeding, these instruments must be able to analyse and draw lessons from the data gathered, either as field or greenhouse experts. For photographs obtained with remote sensing devices or field instruments, image processing has been commonly used in the area of precision agriculture.^[7]

Various types of disease are classified by the following Figure 2;

Picture analysis has been widely used in the literature of agriculture. For example, image processing is used for weed identification and fruit classification, plant disease identification, phenotypic quantification and classification and plant disease symptoms analysis.^[8] Deep learning has recently been used for detection.^[9,10] Mohanty *et al.* In their study, deep learning was used to identify leaf diseases in different plants.^[11] Deep learning is the most popular form of machine learning using hidden layers of artificial neural networks (ANNs).^[12] Classification tasks using semantic characteristics were performed prior to the advent of deep learning patterns.^[12] Deep learning (DL) methods are a branch of machine learning (ML) and were implemented in 1943,^[1] when threshold logic was developed to create computational models that closely mimic human biological pathways. This area of study continues to

grow. Its history can be divided into two phases, from 1943 to 2006 and from 2012 to the present.

Farmers cannot carry out periodic surveys in fields longer than miles at a time to track each plant intensely. In rural areas, too, the shortage of skilled farmers has always been the issue of detecting diseases at the right time. Many developed countries have opened plant clinics for farmers, where they can learn about different diseases and insects. In addition, plant pathologists will identify diseases from samples collected from farmers in these clinics. In addition, agricultural extension officials have also toured the property. Nowadays, fuzzy logic is used to identify plants as diseased or stable plants. If the disease is observed, software (such as fuzzy inference systems) can also be used for classification purposes. Many researchers have used image characteristics (such as colour, form, texture and combinations) to build CBIR systems for various applications. Expand the CBIR method utilising colour functions such as the RGB colour histogram, the GLCM colour histogram and the K-Means colour histogram. Image texture features, such as local binary pattern (LBP) and LBP variance, can be used to build CBIR systems. Researchers use the modulated discrete transform cosine (SADCT) and vector scale function (SIFT) for Canny Acne Detection (CED) as shape characteristics for the design of CBIR systems. The CBIR method, which uses a mixture of colours, shapes and textures, is very successful. Any Gabor colour moments and philtres, autocorrelation maps, BDIP and BVLC moments, Gabor colour moments and texture details, wave and pattern histogram moments, pattern and geometry moments, grey coexistence matrix and Zernike moments, HSV colour moments The mix. Transform description, colour histogram, colour correlation map, grey co-occurrence matrix and Tamura-Hu pair, colour pair and LBP, HSV colour histogram and co-occurrence matrix.^[13]




Figure 2: Different types of disease in plants.

DIFFERENT TYPES OF TOMATO DISEASE

Tomato (*Solanum lycopersicum*) can be grown in almost any well-drained soil. An adequate supply of organic products can improve yields and reduce production issues. Tomatoes and associated vegetables (such as tomatoes, peppers and eggplants) should not be planted several times in the same soil in three years. Ideally, any cover or crop previous to the tomato should be a part of the grass family. Corn is a healthy tomato crop, supplying a significant amount of organic matter and not allowing the growth of pathogenic species that infect tomatoes. The use of approved seeds and plants is advised and can be done whenever possible.

There are many types of diseases in tomatoes, some of the most serious and common diseases will be explained below.

Bacterial wilt

Bacterial soil or bacterial disease is a severe illness caused by *Ralstonia solanacearum* (formerly *Pseudomonas solanacearum*). This type of bacteria can live in the soil for a long time, enter the roots of plants by transplantation, cultivation, or wounds caused by insects and enter the roots by natural wounds. Hot temperatures and humidity contribute to the growth of the disease. Bacteria reproduce quickly in the aquatic plant tubes and fill with mucus. This causes the plant to wilt easily, although the leaves remain green. If the infected stem is transversely sliced, it may look brown and a reduction of yellowish exudate can be seen,^[14-16] Figure 3.

Early Blight

The disease is caused by *Streptomyces tomato* and *Agrobacterium solanum*^[17] and was initially seen as small brown lesions on plants, often on older leaves. The spots are widened and the concentric circles can be seen in the middle of the outbreak, suggestive of bullfighting. The tissue around the stains will turn yellow. If high temperatures and high humidity exist at this time, most leaves will die. Stem diseases are similar to those of trees. If they are close to the soil line (collar), they can occasionally pinch the vine. In the forearm, the lesions attain a large amount, usually occupying nearly the whole forearm. Concentric rings are found in the fruit as well. Infected fruits also break apart. The disease lives in infected soil waste, seeds, volunteer tomato plants and

other Solanaceae hosts (e.g. Irish potatoes, eggplants and black nightshades),^[18-20] Figure 4.

Late blight

Stunting is a potentially severe disease caused by *Phytophthora* in potatoes and tomatoes. Delay is highly harmful in cold, humid climates. The disease is infecting all areas of the vine. The sick spots on the young leaves are tiny and show as dull, water-soaked spots. These leaf spots propagate rapidly and white mould appears along the edges of the infected area on the underside of the leaf. Total defoliation (darkening and shrinking of leaves and stems) may occur within 14 days after the first symptoms begin. Infected tomato fruits have a glossy, black, or olive look that can cover a wide area. Fungal spores disperse between plants and gardens under the influence of rain and wind. Daytime temperatures above 70 degrees Celsius and high humidity are well matched to infection.^[21,22]

Septoria Leaf Spot

This debilitating disease of tomato leaves, roots and roots (the fruit is not infected) is caused by *Septoria lycopersici* fungus. Typically, as the plant starts to bear fruit, the fungus occurs on the lower leaves near the bottom. Many tiny circular spots occur on older leaves, surrounded by dark beige-centered margins. Tiny black spots occur in the middle of the spots that are spore-producing artifacts. The very marked leaves turn yellow and die and fall off the vine. The fungus is more aggressive when the temperature is between 68 and 77° F, the humidity is heavy and the rain or good irrigation



Figure 3: Bacterial wilt.



Figure 4: Early Blight.

moisturises the plants. Falling leaves hinder the growth of the plant, reduce the size and consistency of the fruit and expose the fruit to burning (see below). This fungus does not propagate from the soil but does survive winter crop residues, plant destruction and some tomato-associated wild tomatoes,^[23] Figure 5.

Leaf Mold

The *Passalora fulva* fungus causes leaf mould. This effect was first observed in older leaves near the ground with poor airflow and high humidity. The first signs are a bright green or yellowish spot on the upper surface of the leaves, which expands and becomes noticeable in yellow. In wet conditions, the areas on the lower surface of the leaves are hidden by the velvety and grey growth of the fungus spores. When the infection is severe, the spots merge and the leaves die. Fungi will also infect roots, flowers and fruits. Ripe green fruits may have black leather rot on the tip of the stem.^[24] Fungi live in crop residues and dirt. Spores are dispersed by rain, wind, or instruments. The seeds may have been contaminated. Fungi rely on high relative humidity and high temperature for the propagation of disease,^[25] Figure 6.

Bacterial Spot

The disease is caused by the bacterium *Xanthomonas*, which infects green tomatoes instead of red tomatoes. The pepper is being targeted. The illness is more prevalent during the rainy season. Plant damage includes leaf and fruit spots, resulting in reduced yields, defoliation and sunburn. Symptoms include several



Figure 5: Septoria Leaf Spot.

small, angular to circular, water-soaked spots on the leaves and slightly elevated spots on the fruit. The spots in the leaf may have a yellow halo. The middle sometimes dries up and calls out,^[26] Figure 7.

Tomato Pith Necrosis

Tomato necrosis is typically an early illness that occurs in greenhouses and in high-yield tomato production. When it rains in the spring, however, tomato necrosis will kill tomatoes and occasionally bell peppers in the home gardens. Lung necrosis is caused by a number of soil-borne *Pseudomonas*, including *Pseudomonas crinkle* and *Carnosus carnosus*. These bacteria are called weak pathogens and can invade rapidly growing tomato plants in wet, cold and humid environments, Figure 8.

Anthracnose

Carcinogenicity of tomatoes is caused by the fungus *Colletotrichum coccoides*, which is mainly a pathogen of tomatoes. When the fruit ripens, the signs first appear as small circular jagged patches and then eventually darken. When each region of infection progresses, the lesions will begin to spread over time. In hot, humid and humid climates (with or without rain) the fungus develops salmon-colored spores that arise from the black fungal substance in the centre of the field. These spores are released by the splashing of water,^[27,28] Figure 9.

Southern Blight

Athelia rolfsii infection (synonymous with *Sclerotium rolfsii*) causes the disease. The first symptom is bent palms, which suggest other vision problems. A brown dry rot may occur on the soil line on the base. White fungus growth can be seen with large and small white



Figure 6: Leaf Mold.

sclerotia of mustard seed. Stem lesions grow quickly, spread around the base, causing abrupt and lasting wilting of all antennae. Typically the lesion is concealed by a white fungus. Fungi will also infect fruit where they touch the earth. Fungi have been able to live in soil and plant litter for many years. It is influenced by humidity and high-temperature conditions, Figure 10.

Growth Cracks

When the environmental conditions (heavy rain or irrigation after a drought) promote fast growth during the ripening season, the tomatoes can burst. Any cracks can be deep, allowing rotting species to invade the fruit and cause the fruit to rot,^[29] Figure 11.



Figure 7: Bacterial Spot.



Figure 8: Tomato Pith Necrosis.

DIFFERENT TYPES OF DISEASE IN POTATO PLANTS

Black scurf and Rhizoctonia canker

Rhizobium infection causes the death of potato plants. Potato tubers covered with fungal fruit bodies. Potato tubers covered with fungal fruit bodies.

Symptom: The surface of the tuber is smooth, irregularly coloured, black, or dark brown fungal; the tuber may be deformed; the reddish-brown to black depression of the bean sprouts; the lesions may constrict the main stem, causing the leaves to curl and turn grey, Figure 12.

Potato Early Blight

Symptoms: Lesions with black edges can create concentric circles of raised and sunk tissue on leaves and stems; the lesions are initially circular but become angular; the leaves are necrotic but still attached to the



Figure 9: Anthracnose.



Figure 10: Southern Blight.

plant; the tubers are dark and dry. Lesions have a look like leather or cork with a watery yellowish-green tip.

Cause: Fungus

Note: The emergence of diseases is conducive to the cycle of wet and dry conditions, as well as periods of high humidity and wet leaves, Figure 13.

Flea beetles

Symptoms: Small holes or pits on the leaves give the leaves a characteristic “diffuse” appearance; young plants and seedlings are particularly sensitive; plant

growth may be reduced; if the damage is severe, the plant may be killed; The pest causing damage is a small black beetle (1.5-3.0 mm) that jumps when disturbed; the beetle usually has a shiny appearance.

Cause: Insects.

Comment: Young plants are more susceptible to damage by flea beetles than older plants; older plants can tolerate infestation; flea beetles may overwinter in neighboring weed species, plant debris, or soil; insects can experience it within a year. The second or third generation, Figure 14.



Figure 11: Growth Cracks.

Potato Late Blight

Symptoms: In wet conditions, at the edge of the lesion on the underside of the leaf, irregularly developed brown lesions scatter across the leaf and white sticky spores develop on the ground. When it is dry, the lesion turns dry and dark brown and the tissue falls. Dark green to brown lesions on water-soaked stems often display a distinctive white spore formation; later, the leaves and petioles rot entirely through infection; badly infected plants will have a clear, mildly sweet odour; rough red-brown tuber lesions expand to tissue in the middle of a few centimetres; the presence of the lesion can be mildly bloated and sometimes contribute to secondary lesions.^[30,31] Cause: Oomycete

Comment: Diseases can remain in the soil for months or even years; warm and cool temperatures favour the growth of disease; the major cause of disease transmission is tuber infection,^[32] Figure 15.

Potato leaf roll

Symptoms: The young leaves are twisted, yellow, or pink. The lower leaves are leathery in texture and roll upward; there may be a necrotic network in the tissue



Figure 12: Black scurf and Rhizoctonia canker.



Figure 13: Potato Early Blight.

of the tuber of the blood vessel; the plant has vertical growth and growth may be blocked.^[33] Cause: Virus

Comment: Spread by several aphids; infected tubers and volunteer potato plants are a source of inoculation for the virus, Figure 16.



Figure 14: Flea beetles.



Figure 15: Potato Late Blight.

DIFFERENT TECHNIQUES FOR DETECTION OF TOMATO DISEASE

Deep Learning with Convolution Neural Network (CNN)

Much like Le Chun and so on. Deep learning is a way of reflecting learning in her research. Here, representation learning means that the algorithm can find the best way to replicate the results. This representation is discovered by the algorithm by optimization rather than semantic characteristics. No practical engineering is expected during this learning process. All the products are sold immediately. Mathematically, deep learning is an artificial neural network with unknown layers. Figure 17a displays a neural network of two hidden




Figure 16: Potato leaf roll.




Figure 17 a): The deep neural network, b). Schematic representation of a perceptron.

layers and Figure 1b displays a schematic diagram of the perceptron. Perceptrons have been inspired by live neuronal cells. -- sensor has several inputs (only 5 inputs are seen in Figure 17b) and trigger functions. (1) The mathematical formula for the neuron is given. The activation function makes the neuron's response non-linear, while the network without the activation function is a linear combination of inputs. There are several roles of activation in the literature, such as type S, tanh and ReLU. ReLU is one of the most commonly used stimulus functions and can be trained more quickly.

The weights for each perceptron are calibrated at the beginning of the preparation. The variance from the Gaussian sample can be initialised. At each point of the iteration, all training data will be distributed across the network. As supervised instruction, the deficit between the real field and the grid output will be determined. Loss is the reference to the algorithm for optimization. The optimization algorithm updates the weights on the basis of this failure. The stochastic gradient descent (SGD) algorithm is one of the most commonly used optimization algorithms. In brief, SGD minimises repeated loss on the basis of gradient update processes. There is a special network architecture called the Convolutional Neural Network (CNN) for images such as relatively large, high-dimensional imagery. CNN was used for the first time to find handwritten numbers in documents.^[34] CNN consists of a coherent layer, a focus layer, an activation layer, an escape layer and a totally connected layer. The adhesive layer stores the influence of the philtre or core in combination with the previous layer. These philtres or cores are composed of weights and biases. The aim of the optimization feature is to build certain cores that reflect error-free results. The fused layer is used for downsampling to reduce the size of the neurons and to minimise over-placement. The most commonly played form of billiards is the one with the highest volume that earns the most value in the pool frame. The degree of trigger mode is used to apply non-linearity to the network. The normal stimulus is called the ReLU. The abandonment layer is used to prevent overlaying. The packet loss layer inadvertently switches off the neurons on the network. The completely associated degree is used to measure the chance or ranking for the class. The input of the classifier may be the output of the completely connected layer.

Figure 18 DL deployment flowchart: First, collect the dataset and then split it into two parts, usually divided into 80 percent training and 20 percent validation. After that, develop the DL model^[35] from scratch or use transfer learning strategies to get the training/validation graph to demonstrate the value of the model.

Then, output metrics are used to classify the image (specific categories of plant diseases) and eventually, visualisation/mapping techniques are used to detect/locate/classify pictures.

The dominant neural network is a type of deep neural network capable of processing multidimensional data. CNN's aim is to simplify the picture to a conveniently accessible format without sacrificing the functionality required to make accurate predictions. CNN^[36] has a number of platforms, such as AlexNet, GoogleNet, VGGNet, etc. Its production has given rise to a great deal of interest among researchers in various fields of computer science. It has been used in agriculture to identify plant diseases. As seen in Figure 6, the CNN model contains an input layer, a coherent layer, a focus layer, a totally bonded layer and an output layer. Images are used as inputs to correctly identify diseases in plants. Cohesive layers are used to remove the characteristics of the images. The concentration degree measures the characteristic values of the exported characteristics. Depending on the sensitivity of the file, you can add more convergence and blend for more detail. The




Figure 18: DL deployment flowchart.




Figure 19: Plant disease classification through CNN.

completely connected layer takes the output of the previous level and transforms it into a single vector that can be used as input to the next level. Finally, the degree of development classifies plant diseases,^[37] Figure 19.

The support vector machine (SVM) is a learning algorithm focused on reducing the structural risk that is often used for classification and regression problems. It is designed to optimise the classification cap so as to distinguish the two divisions as far as possible. As seen in Figure 13, for the purpose of collecting acceptable data points, SVM has been extended to this region and is referred to as a hyperplane. The carrier shall mark neighbouring points on the aircraft on both sides of the aircraft. In order to distinguish these vectors, a fixed margin should be used, which should be the largest by which the SVM can be efficiently equipped,^[38-40] Figure 20.

Artificial Intelligence

Data mining today is an efficient and versatile tool that can be used to forecast plant diseases. Thus, by integrating data mining principles with image analysis, we can quickly determine which crops are contaminated, distinguish diseases according to different problems and colours caused by diseases and propose alternative therapies depending on the magnitude of diseases.^[41] 4. Content Based Image Retrieval Technique

CBIR infrastructure functions independently from conventional text library systems. Import images stored in a cluster and then use the features of these images to compare them. The image functionality will be exported automatically^[42,43] Figure 21.

CBIR systems,^[44] mainly use colour, form and texture as core features, such that they work on the first level of the lowest level. Traditional systems allow queries from

users taking data like images, while some systems have additional choices for users, such as palette or sketch entry. In the next step, the system compares the query image to the stored images and fits them correctly with their attribute values and these images will be displayed to the user. The key categories of image searches are discussed below.

Colour Retrieval

The method of extracting details from a color-based picture has been clarified in a variety of ways. However, several of the approaches are variations of the general principle of colour recovery. As the images are used for contrast, they are first analysed and then the colour histogram is taken from the image, indicating the colour ratio of and pixel in the image. The histogram colour extracted from the image is contained in the database. The user will determine the colour ratio of the input picture during the quest while determining the colour histogram. Consider an image whose colour histogram is very similar to the image in the question. Swain and Ballard were the first to develop the histogram mapping method now commonly used. The advancement of the related technologies allows the use of CBIR^[45,46] technologies in more complex systems. The latest Swan and Ballard enhancement strategies include composite colour histograms and area-based colour queries. The effect of this technique is an improvement on the prior art.^[47]

Texture Retrieval

Image similarities may also be based on shape, although they can sound futile. Texture equality can be used to differentiate between the colours and the areas of the graphic. The texture similarity is obtained by comparing




Figure 20: Disease recognition and classification using SVM.




Figure 21: CBIR based system architecture.

the query values and the image contained in the database. Consider comparing the dynamic brightness parameters of two pixel pairs of images. Calculate the value on the basis of the scale, contrast, position and periodicity of the texture analysis. Gabor philtres and fractals are another tools for evaluating image textures. Complete a texture contrast by uploading a query image or choosing textures from the palette. The device will then take into account images whose texture dimensions complement the query image very well.

Shape Retrieval

The third technique is to use type retrieval to extract information from the file. The basic prerequisite for the recovery of image characteristics is by formatting, which is the basic stage of formatting technology. In comparison to texture, the form is a well-defined process. There is a lot of evidence that certain objects in existence are mainly identified by their forms. Many characteristics of the shape of the object (regardless of the size) are processed for each image contained in the database. The question is addressed by manipulating the corresponding properties of the query image and collecting certain images whose capabilities closely equal the query image. Two essential form characteristics are widely used, such as universal characteristics (such as aspect ratio), invariant rounding and local moments and characteristics (such as the set of continuous boundary line segments).

CONCLUSION AND FUTURE SCOPE

In this evaluation paper, we focus primarily on artificial intelligence (AI) technology, image processing (IP) technology, deep learning (DL) technology, vector machine technology (SVM), convergent neural network (CNN), content the detailed description of the identification of different types of diseases in tomato plants based on image retrieval technology (CBIR). It also covers the various types of diseases commonly produced in tomato plants.

The last work has been analyzed and described in dots. It can be concluded that among the technologies used in the existing work, the concept of deep learning through the CNN method has achieved the highest accuracy.

This analysis describes how CNN DL, AI, SVM and CBIR tools are used to diagnose tomato disease. In addition, several imaging/map methods are illustrated for the diagnosis of disease symptoms. While much progress has been made in the last three to four years, there are still some holes in the research listed below:

1. In most of the studies (as mentioned in the previous sections), the dataset of PlantVillage is used to test the accuracy and efficiency of each DL model/architecture. While this dataset contains photos of several diseases of plant species, it has a simple/light history. However, the real world must be taken into account in the actual situation.
2. Superspectral / multispectral imagery is a new technique that has been used in many areas of science (as defined in Section 3). Therefore, particularly though signs are not apparent, sophisticated DL software must be used to diagnose plant diseases.
3. More analytical techniques for visualising plant disease spots should be implemented as they will prevent excessive use of fungicides/pesticides/herbicides, thus saving costs.
4. Over time, the magnitude of plant diseases will change. It is also important to improve/modify the DL model so that it can be identified and categorised during the onset period of the disease.
5. The DL model/architecture must be effective under multiple lighting environments so that the data collection does not only display the current world but must also include photographs taken under natural conditions.
6. Thorough research is required to explain the factors influencing plant diseases, such as the form and scale of the data collection, the learning rate, the illumination and so on.

ACKNOWLEDGMENT

I am hereby thankful to the Department of Computer Science and Engineering, Banasthali Vidyapith, for their support and help.

REFERENCES

1. Wu SG, Bao FS, Xu EY, Wang YX, Chang YF. A Leaf Recognition Algorithm for Plant Classification using Probabilistic Neural Network. IEEE 7th International Symposium on Signal Processing and Information Technology. 2007;11-16.
2. Ying G, Miao L, Yuan Y, Zelin H. A Study on the Method of Image Pre-Processing for Recognition of Crop Diseases. International Conference on Advanced Computer Control. IEEE. 2008.
3. Aji AF, Munajat Q, Pratama AP, Kalamullah H, Setiyawan J, Arymurthy AM. Detection of Palm Oil Leaf Disease with Image Processing and Neural Network Classification on Mobile Device. International Journal of Computer Theory and Engineering. 2013;5(3):528-32.
4. Anand HK, Ashwin PRK. Applying image processing technique to detect plant diseases. International Journal of Modern Engineering Research (IJMER). 2012;2(5):3661-4.
5. Babu MSP, Rao BS. Leaves Recognition Using Back Propagation Neural Network-Advice For Pest and Disease Control On Crops. India Kisan Net: Expert Advisory System. 2007.
6. Camargo S.J. An image-processing based algorithm to automatically identify plant disease visual symptoms. Biosystems Engineering. 2009;102(1):9-21.

7. Gulhane VA, Gurjar AA. Detection of diseases on cotton leaves and its possible diagnosis. *International Journal of Image Processing*. 2011;5(5):590-8.
8. Maheswary P, Srivastav N. Retrieving similar image using Colour moment feature detector and Kmeans clustering of remote sensing images. *International Conference on Computer and Electrical Engineering*. 2008;821-4.
9. Abu-Naser SS, Kashkash KA, Fayyad M. Developing an Expert System for Plant Disease Diagnosis. *Journal of Artificial Intelligence*. 2008;1(2):78-85.
10. Maliappis MT, Ferentinos KP, Passam KP, Sideridis AB. Gims: A Web based Greenhouse Intelligent Management System. *World Journal of Agricultural Sciences*. 2008;4(5):640-7.
11. Miller S, Beed F, Harmon C. Plant disease diagnostic capabilities and networks. *Annual Review of Phytopathology*. 2009;47:15-38.
12. Mohanty SP, Hughes D, Salathe M. Using Deep Learning for Image-Based Plant Disease Detection. *arXiv*. 2016;eprint arXiv:1604.03169.
13. Parsons N, Edmondson R, Song Y. Image analysis and statistical modeling for measurement and quality assessment of ornamental horticulture crops in glasshouses. *Biosystems Engineering*. 2009;104(2):161-8.
14. Pawar P, Turkar V, Patil P. Cucumber disease detection using artificial neural network. *International Conference on Inventive Computation Technologies (ICICT)*, Coimbatore. 2016;1-5.
15. Walia R, Oberoi N, Kumar A, Singh G. Digital Fingerprint and Security Aspects in Internet of Things Against Social Engineering using Advanced Digital Forensics. *Test Engineering and Management*. 2020;83(3):4914-20.
16. Polder G, Blok PM, DeVilliers HAC, DerWolf JM, Kamp J. Potato Virus Y Detection in Seed Potatoes using Deep Learning on Hyperspectral Images. *Front Plant Sci*. 2019;10:209.
17. Bhatnagar V, Singh G, Kumar G, Gupta R. Internet of Things in Smart Agriculture: Applications and Open Challenges. *International Journal of Students' Research in Technology and Management*. 2020;8(1):11-7.
18. Albrechtsen F. Statistical texture measures computed from grey level co-occurrence matrices. *Image Processing Laboratory, Department of Informatics, University of oslo*. 2008;5(5).
19. Preethi G, Sornagopal V. MRI image classification using GLCM texture features. *Green Computing Communication and Electrical Engineering (ICGCCEE)*. 2014;1-6.
20. Rathod AN, Tanawal B, Shah V. Image processing techniques for detection of leaf disease. *Int J Adv Res Comput Sci Softw Eng*. 2013;3(11).
21. Glorot X, Bordes A, Bengio Y. Deep sparse rectifier neural networks. *Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics*. 2011;315-23.
22. Simonyan K, Zisserman A. Very deep convolutional networks for large-scale image recognition. In *Proceedings of the International Conference on Learning Representations, San Diego, CA, USA*. 2015.
23. Tzionas P, Papadakis PE, Manolakis D. Plant leaves classification based on morphological features and fuzzy surface selection technique. *5th International Conference on Technology and Automation ICTA'05, Thessaloniki, Greece*. 2005;365-70.
24. Wang D, Vinson R, Holmes M, Seibel G, Bechar A, Nof S, *et al*. Early Detection of Tomato Spotted Wilt Virus by Hyperspectral Imaging and Outlier Removal Auxiliary Classifier Generative Adversarial Nets (OR-AC-GAN). *Sci Rep*. 2019;9(1):4377.
25. Yu H, Li M, Zhang H, Feng J. Color texture moments for content-based image retrieval. *International Conference on Image Processing*. 2003.
26. Yue J, Zhao W, Mao S, Liu H. Spectral-spatial classification of hyperspectral images using deep convolutional neural networks. *Remote Sens Lett*. 2015;6(6):468-77.
27. Zeiler MD, Fergus R. Visualizing and Understanding Convolutional Networks. *European Conference on Computer Vision*. 2014;818-33.
28. Zhang X, Han L, Dong Y, Shi Y, Huang W, Han L, *et al*. Deep Learning-Based Approach for Automated Yellow Rust Disease Detection from High-Resolution Hyperspectral UAV Images. *Remote Sens*. 2019;11(13):1554.
29. Revathi P, Hemalatha H. Classification of cotton leaf spot diseases using image processing edge detection techniques. *Emerging Trends in Science, Engineering and Technology (INCOSSET), International Conference*. 2012;169-73.
30. Zitnick CL, Dollár P. Edge boxes: Locating object proposals from edges. In *European Conference on Computer Vision*. 2014;391-405.
31. Singh G, Kumar G, Bhatnagar V, Srivastava A, Jyoti K. Pollution management through internet of things: A substantial solution for society. *Humanities and Social Sciences Reviews*. 2019;7(5):1231-7.
32. Gupta D, Singh R. Disease Detection in Plant using Artificial Neural Network. *International Journal for Research in Applied Science and Engineering Technology (IJRASET)*. 2019;7(6):2560-5.
33. He K, Zhang X, Ren S, Sun J. Deep residual learning for image recognition. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition, Las Vegas, NV, USA*. 2016;770-8.
34. Huang J, Kumar SR, Mitra M, Zhu WJ, Zabih R. Image indexing using Colour correlograms. *IEEE Int. Conf. Computer Vision and Pattern Recognition, San Juan, Puerto Rico*. 1997;762-8.
35. Jin X, Jie L, Wang S, Qi H, Li S. Classifying wheat hyperspectral pixels of healthy heads and Fusarium head blight disease using a deep neural network in the wild field. *Remote Sens*. 2018;10(3):395.
36. Kaundal R, Kapoor AS, Raghava PSG. Machine learning techniques in disease forecasting: A case study on rice blast prediction. *BMC Bioinformatics*. 2006;7(1):485.
37. Kho SJ, Manickam S, Malek S, Mosleh M, Dhillon SK. Automated plant identification using artificial neural network and support vector machine. *Frontiers in Life Science*. 2017;10(1):98-107.
38. Krizhevsky SI, Hinton GHE. Image Net Classification with Deep Convolutional Neural Networks. *Advances in Neural Information Processing Systems*. 2012.
39. Kulkarni N. Color thresholding method for image segmentation of natural images. *International Journal of Image, Graphics and Signal Processing*. 2012;4(1):28.
40. Singh G, Dubey OP, Kumar G. A solution to selective forward attack in wireless sensor network. *International Journal of Students' Research in Technology and Management*. 2018;6(4):01-6.
41. LeCun Y, Bengio Y, Hinton G. Deep learning. *Nature*. 2015;521(7553):436-44.
42. Rumpf T, Mahlein A, Steiner U, Oerke E, Plumer HDL. Early detection and classification of plant diseases with support vector machines based on hyperspectral reflectance. *Comput Electron Agr*. 2010;74(1):91-9.
43. Signoroni A, Savardi M, Baronio A, Benini S. Deep Learning Meets Hyperspectral Image Analysis: A Multidisciplinary Review. *Journal of Imaging*. 2019;5(5):52.
44. Sladojevic S, Arsenovic M, Anderla A, Culibrk D, Stefanovic D. Deep Neural Networks Based Recognition of Plant Diseases by Leaf Image Classification. *Computational Intelligence and Neuroscience*. 2016;11.
45. Tellaeche A, Burgos-Artizzu XP, Pajares G, Ribeiro A. A vision-based method for weeds identification through the Bayesian decision theory. *Pattern Recognition*. 2008;41(2):521-30.
46. Gupta R, Singh V. Comparative Analysis of Image Retrieval Techniques in Cyberspace. *International Journal of Students' Research in Technology and Management*. 2020;8(1):1-10.
47. Singh V, Gupta R. Semantics Based Image Retrieval from Cyberspace: A Review Study. *International Journal of Advanced Research in Computer Science*. 2016;7(4):16-21.

Cite this article: Singh G, Yogi KK. A Review on Recognition of Plant Disease using Intelligent Image Retrieval Techniques. *Asian J Biol Life Sci*. 2020;9(3):274-85.